

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
18 April 2002 (18.04.2002)

PCT

(10) International Publication Number  
**WO 02/30981 A1**

(51) International Patent Classification<sup>7</sup>: **C07K 14/705**,  
16/28, C12N 15/12, A01K 67/02, C12P 21/00, C12Q  
1/68, A61K 38/17

(US). **BEASLEY, Ellen, M.**; c/o Celera, 45 West Gude  
Drive, Rockville, MD 20850 (US).

(21) International Application Number: PCT/US01/07832

(74) Agent: **MILLMAN, Robert, A.**; Celera Genomics Corp.,  
45 West Gude Drive C2-4, Rockville, MD 20850 (US).

(22) International Filing Date: 13 March 2001 (13.03.2001)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/684,393 10 October 2000 (10.10.2000) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant: **PE CORPORATION (NY)** [US/US]; Mill-  
man, Robert, A., 761 Main Avenue, Norwalk, CT 06859  
(US).

(72) Inventors: **WEI, Ming-Hui**; c/o Celera, 45 West Gude  
Drive, Rockville, MD 20850 (US). **ZHONG, Wenyan**; c/o  
Celera, 45 West Gude Drive, Rockville, MD 20850 (US).  
**KETCHUM, Karen, A.**; c/o Celera, 45 West Gude Drive,  
Rockville, MD 20850 (US). **DIFRANCESCO, Valentina**;  
c/o Celera, 45 West Gude Drive, Rockville, MD 20850

**Published:**

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: GENE ENCODING A HUMAN G-PROTEIN COUPLED RECEPTOR AND ITS USE

(57) Abstract: The present invention provides amino acid sequences of peptides that are encoded by genes within the Human genome, the GPCR peptides of the present invention. The present invention specifically provides isolated peptide and nucleic acid molecules, methods of identifying orthologs and paralogs of the GPCR peptides and methods of identifying modulators of the GPCR peptides.



WO 02/30981 A1

## GENE ENCODING A HUMAN G-PROTEIN COUPLED RECEPTOR AND ITS USE

**RELATED APPLICATIONS**

5       The present application claims priority to U.S. Serial No. 09/684,393, filed October 10, 2000 (Atty. Docket CL000869).

**FIELD OF THE INVENTION**

10       The present invention is in the field of G-Protein coupled receptors (GPCRs) that are related to the calcium-sensing receptor subfamily, recombinant DNA molecules, and protein production. The present invention specifically provides novel GPCR peptides and proteins and nucleic acid molecules encoding such peptide and protein molecules, all of which are useful in the development of human therapeutics and diagnostic compositions and methods.

15       **BACKGROUND OF THE INVENTION**

G-protein coupled receptors

20       G-protein coupled receptors (GPCRs) constitute a major class of proteins responsible for transducing a signal within a cell. GPCRs have three structural domains: an amino terminal extracellular domain, a transmembrane domain containing seven transmembrane segments, three extracellular loops, and three intracellular loops, and a carboxy terminal intracellular domain. Upon binding of a ligand to an extracellular portion of a GPCR, a signal is transduced within the cell that results in a change in a biological or physiological property of the cell. GPCRs, along with G-proteins and effectors (intracellular enzymes and channels modulated by G-proteins), are the components of a modular signaling system that connects the state of intracellular second  
25       messengers to extracellular inputs.

30       GPCR genes and gene-products are potential causative agents of disease (Spiegel *et al.*, *J. Clin. Invest.* 92:1119-1125 (1993); McKusick *et al.*, *J. Med. Genet.* 30:1-26 (1993)). Specific defects in the rhodopsin gene and the V2 vasopressin receptor gene have been shown to cause various forms of retinitis pigmentosum (Nathans *et al.*, *Annu. Rev. Genet.* 26:403-424(1992)), and nephrogenic diabetes insipidus (Holtzman *et al.*, *Hum. Mol. Genet.* 2:1201-1204 (1993)). These receptors are of critical importance to both the central nervous system and peripheral

physiological processes. Evolutionary analyses suggest that the ancestor of these proteins originally developed in concert with complex body plans and nervous systems.

The GPCR protein superfamily can be divided into five families: Family I, receptors typified by rhodopsin and the  $\beta_2$ -purinergic receptor and currently represented by over 200 unique members (Dohlman *et al.*, *Annu. Rev. Biochem.* 60:653-688 (1991)); Family II, the parathyroid hormone/calcitonin/secretin receptor family (Juppner *et al.*, *Science* 254:1024-1026 (1991); Lin *et al.*, *Science* 254:1022-1024 (1991)); Family III, the metabotropic glutamate receptor family (Nakanishi, *Science* 258 597:603 (1992)); Family IV, the cAMP receptor family, important in the chemotaxis and development of *D. discoideum* (Klein *et al.*, *Science* 241:1467-1472 (1988)); and Family V, the fungal mating pheromone receptors such as STE2 (Kurjan, *Annu. Rev. Biochem.* 61:1097-1129 (1992)).

There are also a small number of other proteins that present seven putative hydrophobic segments and appear to be unrelated to GPCRs; they have not been shown to couple to G-proteins. *Drosophila* expresses a photoreceptor-specific protein, bride of sevenless (boss), a seven-transmembrane-segment protein that has been extensively studied and does not show evidence of being a GPCR (Hart *et al.*, *Proc. Natl. Acad. Sci. USA* 90:5047-5051 (1993)). The gene *frizzled* (*fz*) in *Drosophila* is also thought to be a protein with seven transmembrane segments. Like boss, *fz* has not been shown to couple to G-proteins (Vinson *et al.*, *Nature* 338:263-264 (1989)).

G proteins represent a family of heterotrimeric proteins composed of  $\alpha$ ,  $\beta$  and  $\gamma$  subunits, that bind guanine nucleotides. These proteins are usually linked to cell surface receptors, e.g., receptors containing seven transmembrane segments. Following ligand binding to the GPCR, a conformational change is transmitted to the G protein, which causes the  $\alpha$ -subunit to exchange a bound GDP molecule for a GTP molecule and to dissociate from the  $\beta\gamma$ -subunits. The GTP-bound form of the  $\alpha$ -subunit typically functions as an effector-modulating moiety, leading to the production of second messengers, such as cAMP (e.g., by activation of adenylyl cyclase), diacylglycerol or inositol phosphates. Greater than 20 different types of  $\alpha$ -subunits are known in humans. These subunits associate with a smaller pool of  $\beta$  and  $\gamma$  subunits. Examples of mammalian G proteins include Gi, Go, Gq, Gs and Gt. G proteins are described extensively in Lodish *et al.*, *Molecular Cell Biology*, (Scientific American Books Inc., New York, N.Y., 1995), the contents of which are incorporated herein by reference. GPCRs, G proteins and G protein-linked effector and second messenger systems have been reviewed in *The G-Protein Linked Receptor Fact Book*, Watson *et al.*, eds., Academic Press (1994).

### Calcium-Sensing Receptors

The protein provided by the present invention is highly homologous to calcium-sensing receptors (CaRs), which are GPCRs. CaRs share extensive sequence similarity with odorant and taste receptors. Both CaRs and odorant receptors may be expressed in epithelia.

5 CaRs form dimers held together by disulfide links. Intermolecular interactions between monomers are thought to be essential for CaR activity.

Calcium is vital to a wide array of physiological processes and therefore it is critical that the concentration of calcium in extracellular fluids be kept within a narrow range. Mutations in CaR that increase or decrease the responsiveness of the CaR receptor to  
10 extracellular calcium concentrations are associated with inherited genetic disorders of calcium homeostasis. Therefore, it is likely that CaR is the main regulator of divalent mineral ion excretion.

CaRs are expressed in, and stimulate proliferation of, fibroblasts, where they are involved in calcium-dependent activation of Src and mitogen-activated kinases in response to  
15 extracellular calcium. CaRs may also be expressed in thyroid glands and are likely involved in the etiology of hyper- and hypocalcemic disorders. Naturally occurring mutations of CaRs are associated with several inherited conditions, including familial hypocalciuric hypercalcemia and neonatal severe hyperparathyroidism.

CaRs are also involved in epithelial differentiation and CaRs may be expressed in  
20 keratinocytes where they likely play an essential role in keratinocyte division and differentiation. Deletion of CaR in knockout mice results in visible alterations of epidermis and reduced levels of loricin, a keratinocyte differentiation marker. Epidermis regeneration is a continuous process that is essential for replacement of skin as well as inner linings of organs such as intestines, kidney ducts, and thyroid. The speed of this process is under tight  
25 control of regulatory factors, many of which are unknown. It is possible that CaR levels are elevated in rapidly dividing skin cells, for example, in keratomas and breast tumors. Antibodies derived against CaRs may be used to detect tumors, and synthetic peptide inhibitors that bind CaRs and block its ability to detect calcium may be used as anti-cancer drugs. Short peptides that mimic the CaR dimerization domain could prevent assembly of  
30 functional CaRs.

For a further review of CaRs, see: Oda *et al.*, *J Biol Chem* 2000 Jan 14;275(2):1183-90; Bikle *et al.*, *J Clin Invest* 1996 Feb 15;97(4):1085-93; McNeil *et al.*, *J Biol Chem* 1998 Jan 9;273(2):1114-20; Bai *et al.*, *Proc Natl Acad Sci U S A* 1999 Mar 16;96(6):2834-9;

Emanuel *et al.*, *Mol Endocrinol* 1996 May;10(5):555-65; and Riccardi *et al.*, *Arch Med Res* 1999 Nov-Dec;30(6):436-48.

### Taste Receptors

Two GPCRs, which have been identified in the apical membranes of rat and mouse  
5 taste cells and are differentially dispersed on the tongue and palate, are putative taste  
receptors. These receptors are targets for studies into gustatory processing. These putative  
taste receptors show extensive sequence similarity to calcium-sensing receptors.

For a further review of putative taste receptors, see Smith *et al.*, *Curr Biol* 1999 Jun  
17, 9(12): R453-5.

### Aminergic GPCRs

One family of the GPCRS, Family II, contains receptors for acetylcholine,  
catecholamine, and indoleamine ligands (hereafter referred to as biogenic amines). The  
biogenic amine receptors (aminergic GPCRs) represent a large group of GPCRs that share a  
common evolutionary ancestor and which are present in both vertebrate (deuterostome), and  
15 invertebrate (protostome) lineages. This family of GPCRs includes, but is not limited to the  
5-HT-like, the dopamine-like, the acetylcholine-like, the adrenaline-like and the melatonin-  
like GPCRs.

### Dopamine receptors

The understanding of the dopaminergic system relevance in brain function and disease  
20 developed several decades ago from three diverse observations following drug treatments.  
These were the observations that dopamine replacement therapy improved Parkinson's disease  
symptoms, depletion of dopamine and other catecholamines by reserpine caused depression and  
antipsychotic drugs blocked dopamine receptors. The finding that the dopamine receptor  
binding affinities of typical antipsychotic drugs correlate with their clinical potency led to the  
25 dopamine overactivity hypothesis of schizophrenia (Snyder, S.H., *Am J Psychiatry* 133, 197-202  
(1976); Seeman, P. and Lee, T., *Science* 188, 1217-9 (1975)). Today, dopamine receptors are  
crucial targets in the pharmacological therapy of schizophrenia, Parkinson's disease, Tourette's  
syndrome, tardive dyskinesia and Huntington's disease. The dopaminergic system includes the  
nigrostriatal, mesocorticolimbic and tuberoinfundibular pathways. The nigrostriatal pathway is  
30 part of the striatal motor system and its degeneration leads to Parkinson's disease; the  
mesocorticolimbic pathway plays a key role in reinforcement and in emotional expression and is

the desired site of action of antipsychotic drugs; the tuberoinfundibular pathways regulates prolactin secretion from the pituitary.

Dopamine receptors are members of the G protein coupled receptor superfamily, a large group proteins that share a seven helical membrane-spanning structure and transduce signals through coupling to heterotrimeric guanine nucleotide-binding regulatory proteins (G proteins). Dopamine receptors are classified into subfamilies: D1-like (D1 and D5) and D2-like (D2, D3 and D4) based on their different ligand binding profiles, signal transduction properties, sequence homologies and genomic organizations (Civelli, O., Bunzow, J.R. and Grandy, D.K., *Annu Rev Pharmacol Toxicol* 33, 281-307 (1993)). The D1-like receptors, D1 and D5, stimulate cAMP synthesis through coupling with Gs-like proteins and their genes do not contain introns within their protein coding regions. On the other hand, the D2-like receptors, D2, D3 and D4, inhibit cAMP synthesis through their interaction with Gi-like proteins and share a similar genomic organization which includes introns within their protein coding regions.

#### Serotonin receptors

Serotonin (5-Hydroxytryptamine; 5-HT) was first isolated from blood serum, where it was shown to promote vasoconstriction (Rapport, M.M., Green, A.A. and Page, I.H., *J Biol Chem* 176, 1243-1251 (1948). Interest on a possible relationship between 5-HT and psychiatric disease was spurred by the observations that hallucinogens such as LSD and psilocybin inhibit the actions of 5-HT on smooth muscle preparations (Gaddum, J.H. and Hameed, K.A., *Br J Pharmacol* 9, 240-248 (1954)). This observation lead to the hypothesis that brain 5-HT activity might be altered in psychiatric disorders (Wooley, D.W. and Shaw, E., *Proc Natl Acad Sci US A* 40, 228-231 (1954); Gaddum, J.H. and Picarelli, Z.P., *Br J Pharmacol* 12, 323-328 (1957)). This hypothesis was strengthened by the introduction of tricyclic antidepressants and monoamine oxidase inhibitors for the treatment of major depression and the observation that those drugs affected noradrenaline and 5-HT metabolism. Today, drugs acting on the serotonergic system have been proved to be effective in the pharmacotherapy of psychiatric diseases such as depression, schizophrenia, obsessive-compulsive disorder, panic disorder, generalized anxiety disorder and social phobia as well as migraine, vomiting induced by cancer chemotherapy and gastric motility disorders.

Serotonin receptors represent a very large and diverse family of neurotransmitter receptors. To date thirteen 5-HT receptor proteins coupled to G proteins plus one ligand-gated ion channel receptor (5-HT3) have been described in mammals. This receptor diversity is thought to reflect serotonin's ancient origin as a neurotransmitter and a hormone as well as the

many different roles of 5-HT in mammals. The 5-HT receptors have been classified into seven subfamilies or groups according to their different ligand-binding affinity profiles, molecular structure and intracellular transduction mechanisms (Hoyer, D. et al., *Pharmacol. Rev.* 46, 157-203 (1994)).

## 5        Adrenergic GPCRs

The adrenergic receptors comprise one of the largest and most extensively characterized families within the G-protein coupled receptor "superfamily". This superfamily includes not only adrenergic receptors, but also muscarinic, cholinergic, dopaminergic, serotonergic, and histaminergic receptors. Numerous peptide receptors include glucagon, somatostatin, and vasopressin receptors, as well as sensory receptors for vision (rhodopsin), taste, and olfaction, also belong to this growing family. Despite the diversity of signalling molecules, G-protein coupled receptors all possess a similar overall primary structure, characterized by 7 putative membrane-spanning .alpha. helices (Probst et al., 1992). In the most basic sense, the adrenergic receptors are the physiological sites of action of the catecholamines, epinephrine and norepinephrine. Adrenergic receptors were initially classified as either .alpha. or .beta. by Ahlquist, who demonstrated that the order of potency for a series of agonists to evoke a physiological response was distinctly different at the 2 receptor subtypes (Ahlquist, 1948). Functionally, .alpha. adrenergic receptors were shown to control vasoconstriction, pupil dilation and uterine inhibition, while .beta. adrenergic receptors were implicated in vasorelaxation, myocardial stimulation and bronchodilation (Regan et al., 1990). Eventually, pharmacologists realized that these responses resulted from activation of several distinct adrenergic receptor subtypes. .beta. adrenergic receptors in the heart were defined as .beta..sub.1, while those in the lung and vasculature were termed .beta..sub.2 (Lands et al., 1967).

.alpha. Adrenergic receptors, meanwhile, were first classified based on their anatomical location, as either pre or post-synaptic (.alpha..sub.2 and .alpha..sub.1, respectively) (Langer et al., 1974). This classification scheme was confounded, however, by the presence of .alpha..sub.2 receptors in distinctly non-synaptic locations, such as platelets (Berthelsen and Pettinger, 1977). With the development of radioligand binding techniques, .alpha. adrenergic receptors could be distinguished pharmacologically based on their affinities for the antagonists prazosin or yohimbine (Stark, 1981). Definitive evidence for adrenergic receptor subtypes, however, awaited purification and molecular cloning of adrenergic receptor subtypes. In 1986, the genes for the hamster .beta..sub.2 (Dickson et al.,

1986) and turkey  $\beta_1$  adrenergic receptors (Yarden et al., 1986) were cloned and sequenced. Hydropathy analysis revealed that these proteins contain 7 hydrophobic domains similar to rhodopsin, the receptor for light. Since that time the adrenergic receptor family has expanded to include 3 subtypes of  $\beta$  receptors (Emorine et al., 1989), 3 subtypes of  $\alpha_1$  receptors (Schwinn et al., 1990), and 3 distinct types of  $\beta_2$  receptors (Lomasney et al., 1990).

The cloning, sequencing and expression of  $\alpha$  receptor subtypes from animal tissues has led to the subclassification of the  $\alpha_1$  receptors into  $\alpha_{1d}$  (formerly known as  $\alpha_{1a}$  or  $\alpha_{1a/1d}$ ),  $\alpha_{1b}$  and  $\alpha_{1c}$  (formerly known as  $\alpha_{1c}$ ) subtypes. Each  $\alpha_1$  receptor subtype exhibits its own pharmacologic and tissue specificities. The designation " $\alpha_{1a}$ " is the appellation recently approved by the IUPHAR Nomenclature Committee for the previously designated " $\alpha_{1c}$ " cloned subtype as outlined in the 1995 Receptor and Ion Channel Nomenclature Supplement (Watson and Girdlestone, 1995). The designation  $\alpha_{1a}$  is used throughout this application to refer to this subtype. At the same time, the receptor formerly designated  $\alpha_{1a}$  was renamed  $\alpha_{1d}$ . The new nomenclature is used throughout this application. Stable cell lines expressing these  $\alpha_1$  receptor subtypes are referred to herein; however, these cell lines were deposited with the American Type Culture Collection (ATCC) under the old nomenclature. For a review of the classification of  $\alpha_1$  adrenoceptor subtypes, see, Martin C. Michel, et al., Naunyn-Schmiedeberg's Arch. Pharmacol. (1995) 352:1-10.

The differences in the  $\alpha$  adrenergic receptor subtypes have relevance in pathophysiologic conditions. Benign prostatic hyperplasia, also known as benign prostatic hypertrophy or BPH, is an illness typically affecting men over fifty years of age, increasing in severity with increasing age. The symptoms of the condition include, but are not limited to, increased difficulty in urination and sexual dysfunction. These symptoms are induced by enlargement, or hyperplasia, of the prostate gland. As the prostate increases in size, it impinges on free-flow of fluids through the male urethra. Concomitantly, the increased noradrenergic innervation of the enlarged prostate leads to an increased adrenergic tone of the bladder neck and urethra, further restricting the flow of urine through the urethra.

The  $\alpha_2$  receptors appear to have diverged rather early from either  $\beta$  or  $\alpha_1$  receptors. The  $\alpha_2$  receptors have been broken down into 3 molecularly distinct subtypes termed  $\alpha_2C_2$ ,  $\alpha_2C_4$ , and  $\alpha_2C_{10}$  based on their chromosomal location. These subtypes appear to correspond to the pharmacologically defined  $\alpha_{2B}$ ,  $\alpha_{2C}$ , and  $\alpha_{2A}$  subtypes, respectively (Bylund et



al., 1992). While all the receptors of the adrenergic type are recognized by epinephrine, they are pharmacologically distinct and are encoded by separate genes. These receptors are generally coupled to different second messenger pathways that are linked through G-proteins. Among the adrenergic receptors,  $\beta_1$  and  $\beta_2$  receptors activate the adenylate cyclase,  $\alpha_2$  receptors inhibit adenylate cyclase and  $\alpha_1$  receptors activate phospholipase C pathways, stimulating breakdown of polyphosphoinositides (Chung, F. Z. et al., J. Biol. Chem., 263:4052 (1988)).  $\alpha_1$  and  $\alpha_2$  adrenergic receptors differ in their cell activity for drugs.

Issued US patent that disclose the utility of members of this family of proteins include, but are not limited to, 6,063,785 Phthalimido arylpiperazines useful in the treatment of benign prostatic hyperplasia; 6,060,492 Selective  $\beta_3$  adrenergic agonists; 6,057,350 Alpha 1a adrenergic receptor antagonists; 6,046,192 Phenylethanolaminotetralincarboxamide derivatives; 6,046,183 Method of synergistic treatment for benign prostatic hyperplasia; 6,043,253 Fused piperidine substituted arylsulfonamides as  $\beta_3$ -agonists; 6,043,224 Compositions and methods for treatment of neurological disorders and neurodegenerative diseases; 6,037,354 Alpha 1a adrenergic receptor antagonists; 6,034,106 Oxadiazole benzenesulfonamides as selective  $\beta_3$  Agonist for the treatment of Diabetes and Obesity; 6,011,048 Thiazole benzenesulfonamides as  $\beta_3$  agonists for treatment of diabetes and obesity; 6,008,361 5,994,506 Adrenergic receptor; 5,994,294 Nitrosated and nitrosylated  $\alpha$ -adrenergic receptor antagonist compounds, compositions and their uses; 5,990,128  $\alpha_1$  specific compounds to treat benign prostatic hyperplasia; 5,977,154 Selective  $\beta_3$  adrenergic agonist; 5,977,115 Alpha 1a adrenergic receptor antagonists; 5,939,443 Selective  $\beta_3$  adrenergic agonists; 5,932,538 Nitrosated and nitrosylated  $\alpha$ -adrenergic receptor antagonist compounds, compositions and their uses; 5,922,722 Alpha 1a adrenergic receptor antagonists 26 5,908,830 and 5,861,309 DNA encoding human alpha 1 adrenergic receptors.

### Purinergic GPCRs

#### Purinoceptor P2Y1

P2 purinoceptors have been broadly classified as P2X receptors which are ATP-gated channels; P2Y receptors, a family of G protein-coupled receptors, and P2Z receptors, which mediate nonselective pores in mast cells. Numerous subtypes have been identified for each of

the P2 receptor classes. P2Y receptors are characterized by their selective responsiveness towards ATP and its analogs. Some respond also to UTP. Based on the recommendation for nomenclature of P2 purinoceptors, the P2Y purinoceptors were numbered in the order of cloning. P2Y1, P2Y2 and P2Y3 have been cloned from a variety of species. P2Y1 responds to both ADP and ATP. Analysis of P2Y receptor subtype expression in human bone and 2 osteoblastic cell lines by RT-PCR showed that all known human P2Y receptor subtypes were expressed: P2Y1, P2Y2, P2Y4, P2Y6, and P2Y7 (Maier et al. 1997). In contrast, analysis of brain-derived cell lines suggested that a selective expression of P2Y receptor subtypes occurs in brain tissue.

Leon et al. generated P2Y1-null mice to define the physiologic role of the P2Y1 receptor. (J. Clin. Invest. 104: 1731-1737(1999)) These mice were viable with no apparent abnormalities affecting their development, survival, reproduction, or morphology of platelets, and the platelet count in these animals was identical to that of wildtype mice. However, platelets from P2Y1-deficient mice were unable to aggregate in response to usual concentrations of ADP and displayed impaired aggregation to other agonists, while high concentrations of ADP induced platelet aggregation without shape change. In addition, ADP-induced inhibition of adenylyl cyclase still occurred, demonstrating the existence of an ADP receptor distinct from P2Y1. P2Y1-null mice had no spontaneous bleeding tendency but were resistant to thromboembolism induced by intravenous injection of ADP or collagen and adrenaline. Hence, the P2Y1 receptor plays an essential role in thrombotic states and represents a potential target for antithrombotic drugs. Somers et al. mapped the P2RY1 gene between flanking markers D3S1279 and D3S1280 at a position 173 to 174 cM from the most telomeric markers on the short arm of chromosome 3. (Genomics 44: 127-130 (1997)).

#### Purinoceptor P2Y2

The chloride ion secretory pathway that is defective in cystic fibrosis (CF) can be bypassed by an alternative pathway for chloride ion transport that is activated by extracellular nucleotides. Accordingly, the P2 receptor that mediates this effect is a therapeutic target for improving chloride secretion in CF patients. Parr et al. reported the sequence and functional expression of a cDNA cloned from human airway epithelial cells that encodes a protein with properties of a P2Y nucleotide receptor. (Proc. Nat. Acad. Sci. 91: 3275-3279 (1994)) The human P2RY2 gene was mapped to chromosome 11q13.5-q14.1.

#### Purinoreceptor P2RY4

The P2RY4 receptor appears to be activated specifically by UTP and UDP, but not by ATP and ADP. Activation of this uridine nucleotide receptor resulted in increased inositol phosphate formation and calcium mobilization. The UNR gene is located on chromosome Xq13.

#### Purinoreceptor P2Y6

Somers et al. mapped the P2RY6 gene to 11q13.5, between polymorphic markers D11S1314 and D11S916, and P2RY2 maps within less than 4 cM of P2RY6. (Genomics 44: 127-130 (1997)) This was the first chromosomal clustering of this gene family to be described.

Adenine and uridine nucleotides, in addition to their well established role in intracellular energy metabolism, phosphorylation, and nucleic acid synthesis, also are important extracellular signaling molecules. P2Y metabotropic receptors are GPCRs that mediate the effects of extracellular nucleotides to regulate a wide variety of physiological processes. At least ten subfamilies of P2Y receptors have been identified. These receptor subfamilies differ greatly in their sequences and in their nucleotide agonist selectivities and efficacies.

It has been demonstrated that the P2Y1 receptors are strongly expressed in the brain, but the P2Y2, P2Y4 and P2Y6 receptors are also present. The localisation of one or more of these subtypes on neurons, on glia cells, on brain vasculature or on ventricle ependymal cells was found by in situ mRNA hybridisation and studies on those cells in culture. The P2Y1 receptors are prominent on neurons. The coupling of certain P2Y receptor subtypes to N-type  $\text{Ca}^{2+}$  channels or to particular  $\text{K}^{+}$  channels was also demonstrated.

It has also been demonstrated that several P2Y receptors mediate potent growth stimulatory effects on smooth muscle cells by stimulating intracellular pathways including Gq-proteins, protein kinase C and tyrosine phosphorylation, leading to increased immediate early gene expression, cell number, DNA and protein synthesis. It has been further demonstrated that P2Y regulation plays a mitogenic role in response to the development of atherosclerosis.

It has further been demonstrated that P2Y receptors play a critical role in cystic fibrosis. The volume and composition of the liquid that lines the airway surface is modulated by active transport of ions across the airway epithelium. This in turn is regulated both by autonomic agonists acting on basolateral receptors and by agonists acting on luminal receptors. Specifically, extracellular nucleotides present in the airway surface liquid act on luminal P2Y receptors to control both  $\text{Cl}^{-}$  secretion and  $\text{Na}^{+}$  absorption. Since nucleotides are released in a regulated manner from airway epithelial cells, it is likely that their control over airway ion

transport forms part of an autocrine regulatory system localised to the luminal surface of airway epithelia. In addition to this physiological role, P2Y receptor agonists have the potential to be of crucial benefit in the treatment of CF, a disorder of epithelial ion transport. The airways of people with CF have defective Cl<sup>-</sup> secretion and abnormally high rates of Na<sup>+</sup> absorption. Since P2Y receptor agonists can regulate both these ion transport pathways they have the potential to pharmacologically bypass the ion transport defects in CF.

GPCRs, particularly members of the calcium-sensing receptor subfamily, are a major target for drug action and development. Accordingly, it is valuable to the field of pharmaceutical development to identify and characterize previously unknown GPCRs. The present invention advances the state of the art by providing a previously unidentified human GPCR.

### SUMMARY OF THE INVENTION

The present invention is based in part on the identification of nucleic acid sequences that encode amino acid sequences of human GPCR peptides and proteins that are related to the calcium-sensing receptor subfamily, allelic variants thereof and other mammalian orthologs thereof. These unique peptide sequences, and nucleic acid sequences that encode these peptides, can be used as models for the development of human therapeutic targets, aid in the identification of therapeutic proteins, and serve as targets for the development of human therapeutic agents.

The proteins of the present inventions are GPCRs that participate in signaling pathways mediated by the calcium-sensing receptor subfamily in cells that express these proteins. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. As used herein, a "signaling pathway" refers to the modulation (e.g., stimulation or inhibition) of a cellular function/activity upon the binding of a ligand to the GPCR protein. Examples of such functions include mobilization of intracellular molecules that participate in a signal transduction pathway, e.g., phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>), inositol 1,4,5-triphosphate (IP<sub>3</sub>) and adenylate cyclase; polarization of the plasma membrane; production or secretion of molecules; alteration in the structure of a cellular component; cell proliferation, e.g., synthesis of DNA; cell migration; cell differentiation; and cell survival

The response mediated by the receptor protein depends on the type of cell it is expressed on. Some information regarding the types of cells that express other members of the subfamily

of GPCRs of the present invention is already known in the art (see references cited in Background and information regarding closest homologous protein provided in Figure 2; Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. ). For example, in some cells, binding of a ligand to the receptor protein may stimulate an activity such as release of compounds, gating of a channel, cellular adhesion, migration, differentiation, etc., through phosphatidylinositol or cyclic AMP metabolism and turnover while in other cells, the binding of the ligand will produce a different result. Regardless of the cellular activity/response modulated by the particular GPCR of the present invention, a skilled artisan will clearly know that the receptor protein is a GPCR and interacts with G proteins to produce one or more secondary signals, in a variety of intracellular signal transduction pathways, e.g., through phosphatidylinositol or cyclic AMP metabolism and turnover, in a cell thus participating in a biological process in the cells or tissues that express the GPCR. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Specifically, a virtual northern blot shows expression in a pooled fetal lung/testis/B-cell sample. In addition, PCR-based tissue screening panels indicate expression in humans in Hela cells and bone marrow.

As used herein, "phosphatidylinositol turnover and metabolism" refers to the molecules involved in the turnover and metabolism of phosphatidylinositol 4,5-bisphosphate ( $\text{PIP}_2$ ) as well as to the activities of these molecules.  $\text{PIP}_2$  is a phospholipid found in the cytosolic leaflet of the plasma membrane. Binding of ligand to the receptor activates, in some cells, the plasma-membrane enzyme phospholipase C that in turn can hydrolyze  $\text{PIP}_2$  to produce 1,2-diacylglycerol (DAG) and inositol 1,4,5-triphosphate ( $\text{IP}_3$ ). Once formed  $\text{IP}_3$  can diffuse to the endoplasmic reticulum surface where it can bind an  $\text{IP}_3$  receptor, e.g., a calcium channel protein containing an  $\text{IP}_3$  binding site.  $\text{IP}_3$  binding can induce opening of the channel, allowing calcium ions to be released into the cytoplasm.  $\text{IP}_3$  can also be phosphorylated by a specific kinase to form inositol 1,3,4,5-tetraphosphate ( $\text{IP}_4$ ), a molecule that can cause calcium entry into the cytoplasm from the extracellular medium.  $\text{IP}_3$  and  $\text{IP}_4$  can subsequently be hydrolyzed very rapidly to the inactive products inositol 1,4-bisphosphate ( $\text{IP}_2$ ) and inositol 1,3,4-triphosphate, respectively. These inactive products can be recycled by the cell to synthesize  $\text{PIP}_2$ . The other second messenger produced by the hydrolysis of  $\text{PIP}_2$ , namely 1,2-diacylglycerol (DAG), remains in the cell membrane where it can serve to activate the enzyme protein kinase C. Protein kinase C is usually found soluble in the cytoplasm of the cell, but upon an increase in the intracellular calcium concentration, this enzyme can move to the plasma membrane where it can

be activated by DAG. The activation of protein kinase C in different cells results in various cellular responses such as the phosphorylation of glycogen synthase, or the phosphorylation of various transcription factors, e.g., NF-kB. The language "phosphatidylinositol activity", as used herein, refers to an activity of PIP<sub>2</sub> or one of its metabolites.

5 Another signaling pathway in which the receptor may participate is the cAMP turnover pathway. As used herein, "cyclic AMP turnover and metabolism" refers to the molecules involved in the turnover and metabolism of cyclic AMP (cAMP) as well as to the activities of these molecules. Cyclic AMP is a second messenger produced in response to ligand-induced stimulation of certain G protein coupled receptors. In the cAMP signaling  
10 pathway, binding of a ligand to a GPCR can lead to the activation of the enzyme adenylyl cyclase, which catalyzes the synthesis of cAMP. The newly synthesized cAMP can in turn activate a cAMP-dependent protein kinase. This activated kinase can phosphorylate a voltage-gated potassium channel protein, or an associated protein, and lead to the inability of the potassium channel to open during an action potential. The inability of the potassium  
15 channel to open results in a decrease in the outward flow of potassium, which normally repolarizes the membrane of a neuron, leading to prolonged membrane depolarization.

By targeting an agent to modulate a GPCR, the signaling activity and biological process mediated by the receptor can be agonized or antagonized in specific cells and tissues. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow,  
20 and a pooled sample of fetal lung, testis, and B-cells. Such agonism and antagonism serves as a basis for modulating a biological activity in a therapeutic context (mammalian therapy) or toxic context (anti-cell therapy, e.g. anti-cancer agent).

## DESCRIPTION OF THE FIGURE SHEETS

25 FIGURE 1 provides the nucleotide sequence of a cDNA molecule, with 5' and 3' UTR regions, which encodes the GPCR of the present invention. (SEQ ID NO:1) In addition, structure and functional information is provided, such as ATG start, stop and tissue distribution, where available, that allows one to readily determine specific uses of inventions based on this molecular sequence. Experimental data as provided in Figure 1 indicates  
30 expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells.

FIGURE 2 provides the predicted amino acid sequence of the GPCR of the present invention. (SEQ ID NO:2) In addition structure and functional information such as protein

family, function, and modification sites is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence.

FIGURE 3 provides genomic sequences that span the gene encoding the GPCR protein of the present invention. (SEQ ID NO:3) In addition structure and functional information, such as intron/exon structure, promoter location, etc., is provided where available, allowing one to readily determine specific uses of inventions based on this molecular sequence. As illustrated in Figure 3, known SNP variations include T406C, T852C, G897A, C1433T, T5845C, and G7028A.

## DETAILED DESCRIPTION OF THE INVENTION

### General Description

The present invention is based on the sequencing of the human genome. During the sequencing and assembly of the human genome, analysis of the sequence information revealed previously unidentified fragments of the human genome that encode peptides that share structural and/or sequence homology to protein/peptide/domains identified and characterized within the art as being a GPCR protein or part of a GPCR protein, that are related to the calcium-sensing receptor subfamily. Utilizing these sequences, additional genomic sequences were assembled and transcript and/or cDNA sequences were isolated and characterized. Based on this analysis, the present invention provides amino acid sequences of human GPCR peptides and proteins that are related to the calcium-sensing receptor subfamily, nucleic acid sequences in the form of transcript sequences, cDNA sequences and/or genomic sequences that encode these GPCR peptides and proteins, nucleic acid variation (allelic information), tissue distribution of expression, and information about the closest art known protein/peptide/domain that has structural or sequence homology to the GPCR of the present invention.

In addition to being previously unknown, the peptides that are provided in the present invention are selected based on their ability to be used for the development of commercially important products and services. Specifically, the present peptides are selected based on homology and/or structural relatedness to known GPCR proteins of the calcium-sensing receptor subfamily and the expression pattern observed. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. The art has clearly established the commercial importance of members of

this family of proteins and proteins that have expression patterns similar to that of the present gene. Some of the more specific features of the peptides of the present invention, and the uses thereof, are described herein, particularly in the Background of the Invention and in the annotation provided in the Figures, and/or are known within the art for each of the known calcium-sensing receptor family or subfamily of GPCR proteins.

### Specific Embodiments

#### Peptide Molecules

The present invention provides nucleic acid sequences that encode protein molecules that have been identified as being members of the GPCR family of proteins and are related to the calcium-sensing receptor subfamily (protein sequences are provided in Figure 2, transcript/cDNA sequences are provided in Figure 1 and genomic sequences are provided in Figure 3). The peptide sequences provided in Figure 2, as well as the obvious variants described herein, particularly allelic variants as identified herein and using the information in Figure 3, will be referred herein as the GPCR peptides of the present invention, GPCR peptides, or peptides/proteins of the present invention.

The present invention provides isolated peptide and protein molecules that consist of, consist essentially of, or comprise the amino acid sequences of the GPCR peptides disclosed in Figure 2, (encoded by the nucleic acid molecule shown in Figure 1, transcript/cDNA sequence, or Figure 3, genomic sequence), as well as all obvious variants of these peptides that are within the art to make and use. Some of these variants are described in detail below.

As used herein, a peptide is said to be "isolated" or "purified" when it is substantially free of cellular material or free of chemical precursors or other chemicals. The peptides of the present invention can be purified to homogeneity or other degrees of purity. The level of purification will be based on the intended use. The critical feature is that the preparation allows for the desired function of the peptide, even if in the presence of considerable amounts of other components (the features of an isolated nucleic acid molecule is discussed below).

In some uses, "substantially free of cellular material" includes preparations of the peptide having less than about 30% (by dry weight) other proteins (i.e., contaminating protein), less than about 20% other proteins, less than about 10% other proteins, or less than about 5% other proteins. When the peptide is recombinantly produced, it can also be substantially free of culture



medium, i.e., culture medium represents less than about 20% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of the peptide in which it is separated from chemical precursors or other chemicals that are involved in its synthesis. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of the GPCR peptide having less than about 30% (by dry weight) chemical precursors or other chemicals, less than about 20% chemical precursors or other chemicals, less than about 10% chemical precursors or other chemicals, or less than about 5% chemical precursors or other chemicals.

The isolated GPCR peptide can be purified from cells that naturally express it, purified from cells that have been altered to express it (recombinant), or synthesized using known protein synthesis methods. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. For example, a nucleic acid molecule encoding the GPCR peptide is cloned into an expression vector, the expression vector introduced into a host cell and the protein expressed in the host cell. The protein can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Many of these techniques are described in detail below.

Accordingly, the present invention provides proteins that consist of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). The amino acid sequence of such a protein is provided in Figure 2. A protein consists of an amino acid sequence when the amino acid sequence is the final amino acid sequence of the protein.

The present invention further provides proteins that consist essentially of the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein consists essentially of an amino acid sequence when such an amino acid sequence is present with only a few additional amino acid residues, for example from about 1 to about 100 or so additional residues, typically from 1 to about 20 additional residues in the final protein.

The present invention further provides proteins that comprise the amino acid sequences provided in Figure 2 (SEQ ID NO:2), for example, proteins encoded by the transcript/cDNA nucleic acid sequences shown in Figure 1 (SEQ ID NO:1) and the genomic sequences provided in Figure 3 (SEQ ID NO:3). A protein comprises an amino acid sequence when the amino acid

sequence is at least part of the final amino acid sequence of the protein. In such a fashion, the protein can be only the peptide or have additional amino acid molecules, such as amino acid residues (contiguous encoded sequence) that are naturally associated with it or heterologous amino acid residues/peptide sequences. Such a protein can have a few additional amino acid residues or can comprise several hundred or more additional amino acids. The preferred classes of proteins that are comprised of the GPCR peptides of the present invention are the naturally occurring mature proteins. A brief description of how various types of these proteins can be made/isolated is provided below.

The GPCR peptides of the present invention can be attached to heterologous sequences to form chimeric or fusion proteins. Such chimeric and fusion proteins comprise a GPCR peptide operatively linked to a heterologous protein having an amino acid sequence not substantially homologous to the GPCR peptide. "Operatively linked" indicates that the GPCR peptide and the heterologous protein are fused in-frame. The heterologous protein can be fused to the N-terminus or C-terminus of the GPCR peptide.

In some uses, the fusion protein does not affect the activity of the GPCR peptide *per se*. For example, the fusion protein can include, but is not limited to, enzymatic fusion proteins, for example beta-galactosidase fusions, yeast two-hybrid GAL fusions, poly-His fusions, MYC-tagged, HI-tagged and Ig fusions. Such fusion proteins, particularly poly-His fusions, can facilitate the purification of recombinant GPCR peptide. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a protein can be increased by using a heterologous signal sequence.

A chimeric or fusion protein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different protein sequences are ligated together in-frame in accordance with conventional techniques. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see Ausubel *et al.*, *Current Protocols in Molecular Biology*, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST protein). A GPCR peptide-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the GPCR peptide.

As mentioned above, the present invention also provides and enables obvious variants of the amino acid sequence of the proteins of the present invention, such as naturally occurring

mature forms of the peptide, allelic/sequence variants of the peptides, non-naturally occurring recombinantly derived variants of the peptides, and orthologs and paralogs of the peptides. Such variants can readily be generated using art-known techniques in the fields of recombinant nucleic acid technology and protein biochemistry. It is understood, however, that variants  
5 exclude any amino acid sequences disclosed prior to the invention.

Such variants can readily be identified/made using molecular techniques and the sequence information disclosed herein. Further, such variants can readily be distinguished from other peptides based on sequence and/or structural homology to the GPCR peptides of the present invention. The degree of homology/identity present will be based primarily on whether  
10 the peptide is a functional variant or non-functional variant, the amount of divergence present in the paralog family and the evolutionary distance between the orthologs.

To determine the percent identity of two amino acid sequences or two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for  
15 optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, 40%, 50%, 60%, 70%, 80%, or 90% or more of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first  
20 sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the  
25 length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity and similarity between two sequences can be accomplished using a mathematical algorithm. (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993;  
30 *Computer Analysis of Sequence Data, Part 1*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453

(1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (Devereux, J., *et al.*, *Nucleic Acids Res.* 12(1):387 (1984)) (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Meyers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences of the present invention can further be used as a "query sequence" to perform a search against sequence databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. (*J. Mol. Biol.* 215:403-10 (1990)). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to the nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to the proteins of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (*Nucleic Acids Res.* 25(17):3389-3402 (1997)). When utilizing BLAST and gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

Full-length pre-processed forms, as well as mature processed forms, of proteins that comprise one of the peptides of the present invention can readily be identified as having complete sequence identity to one of the GPCR peptides of the present invention as well as being encoded by the same genetic locus as the GPCR peptide provided herein. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 1.

Allelic variants of a GPCR peptide can readily be identified as being a human protein having a high degree (significant) of sequence homology/identity to at least a portion of the GPCR peptide as well as being encoded by the same genetic locus as the GPCR peptide provided herein. Genetic locus can readily be determined based on the genomic information provided in Figure 3, such as the genomic sequence mapped to the reference human. As

indicated by the data presented in Figure 3, the map position was determined to be on chromosome 1. As used herein, two proteins (or a region of the proteins) have significant homology when the amino acid sequences are typically at least about 70-80%, 80-90%, and more typically at least about 90-95% or more homologous. A significantly homologous amino acid sequence, according to the present invention, will be encoded by a nucleic acid sequence that will hybridize to a GPCR peptide encoding nucleic acid molecule under stringent conditions as more fully described below.

Figure 3 provides information on SNPs that have been found in a gene encoding the GPCR proteins of the present invention. The following SNPs were found: T406C, T852C, G897A, C1433T, T5845C, and G7028A.

Paralogs of a GPCR peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the GPCR peptide, as being encoded by a gene from humans, and as having similar activity or function. Two proteins will typically be considered paralogs when the amino acid sequences are typically at least about 60% or greater, and more typically at least about 70% or greater homology through a given region or domain. Such paralogs will be encoded by a nucleic acid sequence that will hybridize to a GPCR peptide encoding nucleic acid molecule under moderate to stringent conditions as more fully described below.

Orthologs of a GPCR peptide can readily be identified as having some degree of significant sequence homology/identity to at least a portion of the GPCR peptide as well as being encoded by a gene from another organism. Preferred orthologs will be isolated from mammals, preferably primates, for the development of human therapeutic targets and agents. Such orthologs will be encoded by a nucleic acid sequence that will hybridize to a GPCR peptide encoding nucleic acid molecule under moderate to stringent conditions, as more fully described below, depending on the degree of relatedness of the two organisms yielding the proteins.

Non-naturally occurring variants of the GPCR peptides of the present invention can readily be generated using recombinant techniques. Such variants include, but are not limited to deletions, additions and substitutions in the amino acid sequence of the GPCR peptide. For example, one class of substitutions are conserved amino acid substitution. Such substitutions are those that substitute a given amino acid in a GPCR peptide by another amino acid of like characteristics. Typically seen as conservative substitutions are the replacements, one for another, among the aliphatic amino acids Ala, Val, Leu, and Ile; interchange of the hydroxyl residues Ser and Thr; exchange of the acidic residues Asp and Glu; substitution between the

amide residues Asn and Gln; exchange of the basic residues Lys and Arg; and replacements among the aromatic residues Phe and Tyr. Guidance concerning which amino acid changes are likely to be phenotypically silent are found in Bowie *et al.*, *Science* 247:1306-1310 (1990).

Variant GPCR peptides can be fully functional or can lack function in one or more activities, e.g. ability to bind ligand, ability to bind G-protein, ability to mediate signaling, etc. Fully functional variants typically contain only conservative variation or variation in non-critical residues or in non-critical regions. Figure 2 provides the result of protein analysis that identifies critical domains/regions. Functional variants can also contain substitution of similar amino acids that result in no change or an insignificant change in function. Alternatively, such substitutions may positively or negatively affect function to some degree.

Non-functional variants typically contain one or more non-conservative amino acid substitutions, deletions, insertions, inversions, or truncation or a substitution, insertion, inversion, or deletion in a critical residue or critical region.

Amino acids that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham *et al.*, *Science* 244:1081-1085 (1989)), particularly using the results provided in Figure 2. The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as ligand/effector molecule binding or in assays such as an *in vitro* proliferative activity. Sites that are critical for ligand-receptor binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992); de Vos *et al.*, *Science* 255:306-312 (1992)).

The present invention further provides fragments of the GPCR peptides, in addition to proteins and peptides that comprise and consist of such fragments, particularly those comprising the residues identified in Figure 2. The fragments to which the invention pertains, however, are not to be construed as encompassing fragments that may be disclosed publicly prior to the present invention.

As used herein, a fragment comprises at least 8, 10, 12, 14, 16, or more contiguous amino acid residues from a GPCR peptide. Such fragments can be chosen based on the ability to retain one or more of the biological activities of the GPCR peptide or could be chosen for the ability to perform a function, e.g. ability to bind ligand or effector molecule or act as an immunogen. Particularly important fragments are biologically active fragments, peptides which are, for example, about 8 or more amino acids in length. Such fragments will typically comprise a domain or motif of the GPCR peptide, e.g., active site, a G-protein binding site, a

transmembrane domain or a ligand-binding domain. Further, possible fragments include, but are not limited to, domain or motif containing fragments, soluble peptide fragments, and fragments containing immunogenic structures. Predicted domains and functional sites are readily identifiable by computer programs well-known and readily available to those of skill in the art (e.g., PROSITE analysis). The results of one such analysis are provided in Figure 2.

Polypeptides often contain amino acids other than the 20 amino acids commonly referred to as the 20 naturally occurring amino acids. Further, many amino acids, including the terminal amino acids, may be modified by natural processes, such as processing and other post-translational modifications, or by chemical modification techniques well known in the art. Common modifications that occur naturally in GPCR peptides are described in basic texts, detailed monographs, and the research literature, and they are well known to those of skill in the art (some of these features are identified in Figure 2).

Known modifications include, but are not limited to, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent crosslinks, formation of cystine, formation of pyroglutamate, formylation, gamma carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

Such modifications are well-known to those of skill in the art and have been described in great detail in the scientific literature. Several particularly common modifications, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, for instance, are described in most basic texts, such as *Proteins - Structure and Molecular Properties*, 2nd Ed., T.E. Creighton, W. H. Freeman and Company, New York (1993). Many detailed reviews are available on this subject, such as by Wold, F., *Posttranslational Covalent Modification of Proteins*, B.C. Johnson, Ed., Academic Press, New York 1-12 (1983); Seifter *et al.* (*Meth. Enzymol.* 182: 626-646 (1990)) and Rattan *et al.* (*Ann. N.Y. Acad. Sci.* 663:48-62 (1992)).

Accordingly, the GPCR peptides of the present invention also encompass derivatives or analogs in which a substituted amino acid residue is not one encoded by the genetic code, in which a substituent group is included, in which the mature GPCR peptide is fused with another compound, such as a compound to increase the half-life of the GPCR peptide (for example,

polyethylene glycol), or in which the additional amino acids are fused to the mature GPCR peptide, such as a leader or secretory sequence or a sequence for purification of the mature GPCR peptide or a pro-protein sequence.

## 5        Protein/Peptide Uses

The proteins of the present invention can be used in substantial and specific assays related to the functional information provided in the Figures and Back Ground Section; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its binding  
10 partner or receptor) in biological fluids; and as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state). Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the binding partner so as to develop a system to identify inhibitors of  
15 the binding interaction. Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as commercial products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis  
20 eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

The potential uses of the peptides of the present invention are based primarily on the source of the protein as well as the class/action of the protein. For example, GPCRs isolated from humans and their human/mammalian orthologs serve as targets for identifying agents  
25 for use in mammalian therapeutic applications, e.g. a human drug, particularly in modulating a biological or pathological response in a cell or tissue that expresses the GPCR. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Specifically, a virtual northern blot shows expression in a pooled fetal  
30 lung/testis/B-cell sample. In addition, PCR-based tissue screening panels indicate expression in humans in Hela cells and bone marrow. Approximately 70% of all pharmaceutical agents modulate the activity of a GPCR. A combination of the invertebrate and mammalian ortholog can be used in selective screening methods to find agents specific for invertebrates.



The structural and functional information provided in the Background and Figures provide specific and substantial uses for the molecules of the present invention, particularly in combination with the expression information provided in Figure 1. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Such uses can readily be determined using the information provided herein, that known in the art and routine experimentation.

The proteins of the present invention (including variants and fragments that may have been disclosed prior to the present invention) are useful for biological assays related to GPCRs that are related to members of the calcium-sensing receptor subfamily. Such assays involve any of the known GPCR functions or activities or properties useful for diagnosis and treatment of GPCR-related conditions that are specific for the subfamily of GPCRs that the one of the present invention belongs to, particularly in cells and tissues that express this receptor. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Specifically, a virtual northern blot shows expression in a pooled fetal lung/testis/B-cell sample. In addition, PCR-based tissue screening panels indicate expression in humans in Hela cells and bone marrow.

The proteins of the present invention are also useful in drug screening assays, in cell-based or cell-free systems. Cell-based systems can be native, i.e., cells that normally express the receptor protein, as a biopsy or expanded in cell culture. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. In an alternate embodiment, cell-based assays involve recombinant host cells expressing the receptor protein.

The polypeptides can be used to identify compounds that modulate receptor activity of the protein in its natural state, or an altered form that causes a specific disease or pathology associated with the receptor. Both the GPCRs of the present invention and appropriate variants and fragments can be used in high-throughput screens to assay candidate compounds for the ability to bind to the receptor. These compounds can be further screened against a functional receptor to determine the effect of the compound on the receptor activity. Further, these compounds can be tested in animal or invertebrate systems to determine activity/effectiveness. Compounds can be identified that activate (agonist) or inactivate (antagonist) the receptor to a desired degree.

Further, the proteins of the present invention can be used to screen a compound for the ability to stimulate or inhibit interaction between the receptor protein and a molecule that

normally interacts with the receptor protein, e.g. a ligand or a component of the signal pathway that the receptor protein normally interacts (for example, a G-protein or other interactor involved in cAMP or phosphatidylinositol turnover and/or adenylate cyclase, or phospholipase C activation). Such assays typically include the steps of combining the receptor protein with a candidate compound under conditions that allow the receptor protein, or fragment, to interact with the target molecule, and to detect the formation of a complex between the protein and the target or to detect the biochemical consequence of the interaction with the receptor protein and the target, such as any of the associated effects of signal transduction such as G-protein phosphorylation, cAMP or phosphatidylinositol turnover, and adenylate cyclase or phospholipase C activation.

Candidate compounds include, for example, 1) peptides such as soluble peptides, including Ig-tailed fusion peptides and members of random peptide libraries (see, e.g., Lam *et al.*, *Nature* 354:82-84 (1991); Houghten *et al.*, *Nature* 354:84-86 (1991)) and combinatorial chemistry-derived molecular libraries made of D- and/or L- configuration amino acids; 2) phosphopeptides (e.g., members of random and partially degenerate, directed phosphopeptide libraries, see, e.g., Songyang *et al.*, *Cell* 72:767-778 (1993)); 3) antibodies (e.g., polyclonal, monoclonal, humanized, anti-idiotypic, chimeric, and single chain antibodies as well as Fab, F(ab')<sub>2</sub>, Fab expression library fragments, and epitope-binding fragments of antibodies); and 4) small organic and inorganic molecules (e.g., molecules obtained from combinatorial and natural product libraries).

One candidate compound is a soluble fragment of the receptor that competes for ligand binding. Other candidate compounds include mutant receptors or appropriate fragments containing mutations that affect receptor function and thus compete for ligand. Accordingly, a fragment that competes for ligand, for example with a higher affinity, or a fragment that binds ligand but does not allow release, is encompassed by the invention.

The invention further includes other end point assays to identify compounds that modulate (stimulate or inhibit) receptor activity. The assays typically involve an assay of events in the signal transduction pathway that indicate receptor activity. Thus, a cellular process such as proliferation, the expression of genes that are up- or down-regulated in response to the receptor protein dependent signal cascade, can be assayed. In one embodiment, the regulatory region of such genes can be operably linked to a marker that is easily detectable, such as luciferase.

Any of the biological or biochemical functions mediated by the receptor can be used as an endpoint assay. These include all of the biochemical or biochemical/biological events

described herein, in the references cited herein, incorporated by reference for these endpoint assay targets, and other functions known to those of ordinary skill in the art or that can be readily identified using the information provided in the Figures, particularly Figure 2. Specifically, a biological function of a cell or tissues that expresses the receptor can be assayed. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Specifically, a virtual northern blot shows expression in a pooled fetal lung/testis/B-cell sample. In addition, PCR-based tissue screening panels indicate expression in humans in Hela cells and bone marrow.

Binding and/or activating compounds can also be screened by using chimeric receptor proteins in which the amino terminal extracellular domain, or parts thereof, the entire transmembrane domain or subregions, such as any of the seven transmembrane segments or any of the intracellular or extracellular loops and the carboxy terminal intracellular domain, or parts thereof, can be replaced by heterologous domains or subregions. For example, a G-protein-binding region can be used that interacts with a different G-protein than that which is recognized by the native receptor. Accordingly, a different set of signal transduction components is available as an end-point assay for activation. Alternatively, the entire transmembrane portion or subregions (such as transmembrane segments or intracellular or extracellular loops) can be replaced with the entire transmembrane portion or subregions specific to a host cell that is different from the host cell from which the amino terminal extracellular domain and/or the G-protein-binding region are derived. This allows for assays to be performed in other than the specific host cell from which the receptor is derived. Alternatively, the amino terminal extracellular domain (and/or other ligand-binding regions) could be replaced by a domain (and/or other binding region) binding a different ligand, thus, providing an assay for test compounds that interact with the heterologous amino terminal extracellular domain (or region) but still cause signal transduction. Finally, activation can be detected by a reporter gene containing an easily detectable coding region operably linked to a transcriptional regulatory sequence that is part of the native signal transduction pathway.

The proteins of the present invention are also useful in competition binding assays in methods designed to discover compounds that interact with the receptor. Thus, a compound is exposed to a receptor polypeptide under conditions that allow the compound to bind or to otherwise interact with the polypeptide (Hodgson, Bio/technology, 1992, Sept 10(9);973-80). Soluble receptor polypeptide is also added to the mixture. If the test compound interacts with the soluble receptor polypeptide, it decreases the amount of complex formed or activity from the

receptor target. This type of assay is particularly useful in cases in which compounds are sought that interact with specific regions of the receptor. Thus, the soluble polypeptide that competes with the target receptor region is designed to contain peptide sequences corresponding to the region of interest.

5 To perform cell free drug screening assays, it is sometimes desirable to immobilize either the receptor protein, or fragment, or its target molecule to facilitate separation of complexes from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay.

10 Techniques for immobilizing proteins on matrices can be used in the drug screening assays. In one embodiment, a fusion protein can be provided which adds a domain that allows the protein to be bound to a matrix. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the cell lysates (e.g., <sup>35</sup>S-labeled) and the candidate compound, and the mixture incubated under conditions conducive to complex  
15 formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads are washed to remove any unbound label, and the matrix immobilized and radiolabel determined directly, or in the supernatant after the complexes are dissociated. Alternatively, the complexes can be dissociated from the matrix, separated by SDS-PAGE, and the level of receptor-binding protein found in the bead fraction quantitated from the gel using standard electrophoretic  
20 techniques. For example, either the polypeptide or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin using techniques well known in the art. Alternatively, antibodies reactive with the protein but which do not interfere with binding of the protein to its target molecule can be derivatized to the wells of the plate, and the protein trapped in the wells by antibody conjugation. Preparations of a receptor-binding protein and a candidate  
25 compound are incubated in the receptor protein-presenting wells and the amount of complex trapped in the well can be quantitated. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the receptor protein target molecule, or which are reactive with receptor protein and compete with the target molecule, as well as enzyme-linked  
30 assays which rely on detecting an enzymatic activity associated with the target molecule.

Agents that modulate one of the GPCRs of the present invention can be identified using one or more of the above assays, alone or in combination. It is generally preferable to use a cell-based or cell free system first and then confirm activity in an animal or other model system. Such model systems are well known in the art and can readily be employed in this context.

Modulators of receptor protein activity identified according to these drug screening assays can be used to treat a subject with a disorder mediated by the receptor pathway, by treating cells or tissues that express the GPCR. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. These methods of treatment include the steps of administering a modulator of the GPCR's activity in a pharmaceutical composition to a subject in need of such treatment, the modulator being identified as described herein.

In yet another aspect of the invention, the GPCR proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No. 5,283,317; Zervos et al. (1993) *Cell* 72:223-232; Madura et al. (1993) *J. Biol. Chem.* 268:12046-12054; Bartel et al. (1993) *Biotechniques* 14:920-924; Iwabuchi et al. (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with the GPCR and are involved in GPCR activity. Such GPCR-binding proteins are also likely to be involved in the propagation of signals by the GPCR proteins or GPCR targets as, for example, downstream elements of a GPCR-mediated signaling pathway. Alternatively, such GPCR-binding proteins are likely to be GPCR inhibitors.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a GPCR protein is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a GPCR-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the GPCR protein.

This invention further pertains to novel agents identified by the above-described screening assays. Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a GPCR modulating agent, an antisense GPCR nucleic

acid molecule, a GPCR-specific antibody, or a GPCR-binding partner) can be used in an animal or other model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal or other model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

The GPCR proteins of the present invention are also useful to provide a target for diagnosing a disease or predisposition to disease mediated by the peptide. Accordingly, the invention provides methods for detecting the presence, or levels of, the protein (or encoding mRNA) in a cell, tissue, or organism. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. The method involves contacting a biological sample with a compound capable of interacting with the receptor protein such that the interaction can be detected. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

One agent for detecting a protein in a sample is an antibody capable of selectively binding to protein. A biological sample includes tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject.

The peptides of the present invention also provide targets for diagnosing active protein activity, disease, or predisposition to disease, in a patient having a variant peptide, particularly activities and conditions that are known for other members of the family of proteins to which the present one belongs. Thus, the peptide can be isolated from a biological sample and assayed for the presence of a genetic mutation that results in aberrant peptide. This includes amino acid substitution, deletion, insertion, rearrangement, (as the result of aberrant splicing events), and inappropriate post-translational modification. Analytic methods include altered electrophoretic mobility, altered tryptic peptide digest, altered receptor activity in cell-based or cell-free assay, alteration in ligand or antibody-binding pattern, altered isoelectric point, direct amino acid sequencing, and any other of the known assay techniques useful for detecting mutations in a protein. Such an assay can be provided in a single detection format or a multi-detection format such as an antibody chip array.

*In vitro* techniques for detection of peptide include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence using a detection reagent, such as an antibody or protein binding agent. Alternatively, the peptide can be detected *in vivo* in a subject by introducing into the subject a labeled anti-peptide antibody or other types of detection agent. For example, the antibody can be labeled with a radioactive

marker whose presence and location in a subject can be detected by standard imaging techniques. Particularly useful are methods that detect the allelic variant of a peptide expressed in a subject and methods which detect fragments of a peptide in a sample.

The peptides are also useful in pharmacogenomic analysis. Pharmacogenomics deal with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Eichelbaum, M. (*Clin. Exp. Pharmacol. Physiol.* 23(10-11):983-985 (1996)), and Linder, M.W. (*Clin. Chem.* 43(2):254-266 (1997)). The clinical outcomes of these variations result in severe toxicity of therapeutic drugs in certain individuals or therapeutic failure of drugs in certain individuals as a result of individual variation in metabolism. Thus, the genotype of the individual can determine the way a therapeutic compound acts on the body or the way the body metabolizes the compound. Further, the activity of drug metabolizing enzymes effects both the intensity and duration of drug action. Thus, the pharmacogenomics of the individual permit the selection of effective compounds and effective dosages of such compounds for prophylactic or therapeutic treatment based on the individual's genotype. The discovery of genetic polymorphisms in some drug metabolizing enzymes has explained why some patients do not obtain the expected drug effects, show an exaggerated drug effect, or experience serious toxicity from standard drug dosages. Polymorphisms can be expressed in the phenotype of the extensive metabolizer and the phenotype of the poor metabolizer. Accordingly, genetic polymorphism may lead to allelic protein variants of the receptor protein in which one or more of the receptor functions in one population is different from those in another population. The peptides thus allow a target to ascertain a genetic predisposition that can affect treatment modality. Thus, in a ligand-based treatment, polymorphism may give rise to amino terminal extracellular domains and/or other ligand-binding regions that are more or less active in ligand binding, and receptor activation. Accordingly, ligand dosage would necessarily be modified to maximize the therapeutic effect within a given population containing a polymorphism. As an alternative to genotyping, specific polymorphic peptides could be identified.

The peptides are also useful for treating a disorder characterized by an absence of, inappropriate, or unwanted expression of the protein. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Accordingly, methods for treatment include the use of the GPCR protein or fragments.

### Antibodies

The invention also provides antibodies that selectively bind to one of the peptides of the present invention, a protein comprising such a peptide, as well as variants and fragments thereof. As used herein, an antibody selectively binds a target peptide when it binds the target peptide and does not significantly bind to unrelated proteins. An antibody is still considered to selectively bind a peptide even if it also binds to other proteins that are not substantially homologous with the target peptide so long as such proteins share homology with a fragment or domain of the peptide target of the antibody. In this case, it would be understood that antibody binding to the peptide is still selective despite some degree of cross-reactivity.

As used herein, an antibody is defined in terms consistent with that recognized within the art: they are multi-subunit proteins produced by a mammalian organism in response to an antigen challenge. The antibodies of the present invention include polyclonal antibodies and monoclonal antibodies, as well as fragments of such antibodies, including, but not limited to, Fab or F(ab')<sub>2</sub>, and Fv fragments.

Many methods are known for generating and/or identifying antibodies to a given target peptide. Several such methods are described by Harlow, Antibodies, Cold Spring Harbor Press, (1989).

In general, to generate antibodies, an isolated peptide is used as an immunogen and is administered to a mammalian organism, such as a rat, rabbit or mouse. The full-length protein, an antigenic peptide fragment or a fusion protein can be used. Particularly important fragments are those covering functional domains, such as the domains identified in Figure 2, and domain of sequence homology or divergence amongst the family, such as those that can readily be identified using protein alignment methods and as presented in the Figures.

Antibodies are preferably prepared from regions or discrete fragments of the GPCR proteins. Antibodies can be prepared from any region of the peptide as described herein. However, preferred regions will include those involved in function/activity and/or receptor/binding partner interaction. Figure 2 can be used to identify particularly important regions while sequence alignment can be used to identify conserved and unique sequence fragments.

An antigenic fragment will typically comprise at least 8 contiguous amino acid residues. The antigenic peptide can comprise, however, at least 10, 12, 14, 16 or more amino acid residues. Such fragments can be selected on a physical property, such as fragments correspond



to regions that are located on the surface of the protein, e.g., hydrophilic regions or can be selected based on sequence uniqueness (see Figure 2).

Detection on an antibody of the present invention can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$  or  $^3\text{H}$ .

#### Antibody Uses

The antibodies can be used to isolate one of the proteins of the present invention by standard techniques, such as affinity chromatography or immunoprecipitation. The antibodies can facilitate the purification of the natural protein from cells and recombinantly produced protein expressed in host cells. In addition, such antibodies are useful to detect the presence of one of the proteins of the present invention in cells or tissues to determine the pattern of expression of the protein among various tissues in an organism and over the course of normal development. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Specifically, a virtual northern blot shows expression in a pooled fetal lung/testis/B-cell sample. In addition, PCR-based tissue screening panels indicate expression in humans in Hela cells and bone marrow. Further, such antibodies can be used to detect protein *in situ*, *in vitro*, or in a cell lysate or supernatant in order to evaluate the abundance and pattern of expression. Also, such antibodies can be used to assess abnormal tissue distribution or abnormal expression during development or progression of a biological condition. Antibody detection of circulating fragments of the full length protein can be used to identify turnover.

Further, the antibodies can be used to assess expression in disease states such as in active stages of the disease or in an individual with a predisposition toward disease related to the protein's function. When a disorder is caused by an inappropriate tissue distribution,

developmental expression, level of expression of the protein, or expressed/processed form, the antibody can be prepared against the normal protein. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. If a disorder is characterized by a specific mutation in the protein, antibodies specific for this mutant protein can be used to assay for the presence of the specific mutant protein.

The antibodies can also be used to assess normal and aberrant subcellular localization of cells in the various tissues in an organism. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells.

The diagnostic uses can be applied, not only in genetic testing, but also in monitoring a treatment modality. Accordingly, where treatment is ultimately aimed at correcting expression level or the presence of aberrant sequence and aberrant tissue distribution or developmental expression, antibodies directed against the protein or relevant fragments can be used to monitor therapeutic efficacy.

Additionally, antibodies are useful in pharmacogenomic analysis. Thus, antibodies prepared against polymorphic proteins can be used to identify individuals that require modified treatment modalities. The antibodies are also useful as diagnostic tools as an immunological marker for aberrant protein analyzed by electrophoretic mobility, isoelectric point, tryptic peptide digest, and other physical assays known to those in the art.

The antibodies are also useful for tissue typing. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Thus, where a specific protein has been correlated with expression in a specific tissue, antibodies that are specific for this protein can be used to identify a tissue type.

The antibodies are also useful for inhibiting protein function, for example, blocking the binding of the GPCR peptide to a binding partner such as a ligand. These uses can also be applied in a therapeutic context in which treatment involves inhibiting the protein's function. An antibody can be used, for example, to block binding, thus modulating (agonizing or antagonizing) the peptides activity. Antibodies can be prepared against specific fragments containing sites required for function or against intact protein that is associated with a cell or cell membrane. See Figure 2 for structural information relating to the proteins of the present invention.

The invention also encompasses kits for using antibodies to detect the presence of a protein in a biological sample. The kit can comprise antibodies such as a labeled or labelable antibody and a compound or agent for detecting protein in a biological sample; means for determining the amount of protein in the sample; means for comparing the amount of protein in

the sample with a standard; and instructions for use. Such a kit can be supplied to detect a single protein or epitope or can be configured to detect one of a multitude of epitopes, such as in an antibody detection array. Arrays are described in detail below for nucleic acid arrays and similar methods have been developed for antibody arrays.

5

### Nucleic Acid Molecules

The present invention further provides isolated nucleic acid molecules that encode a GPCR peptide or protein of the present invention (cDNA, transcript and genomic sequence). Such nucleic acid molecules will consist of, consist essentially of, or comprise a nucleotide  
10 sequence that encodes one of the GPCR peptides of the present invention, an allelic variant thereof, or an ortholog or paralog thereof.

As used herein, an "isolated" nucleic acid molecule is one that is separated from other nucleic acid present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. However, there can be some flanking nucleotide sequences, for example up to  
15 about 5KB, 4KB, 3KB, 2KB, or 1KB or less, particularly contiguous peptide encoding sequences and peptide encoding sequences within the same gene but separated by introns in the genomic sequence. The important point is that the nucleic acid is isolated from remote and unimportant flanking sequences such that it can be subjected to the specific manipulations  
20 described herein such as recombinant expression, preparation of probes and primers, and other uses specific to the nucleic acid sequences.

Moreover, an "isolated" nucleic acid molecule, such as a transcript/cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by  
25 recombinant techniques, or chemical precursors or other chemicals when chemically synthesized. However, the nucleic acid molecule can be fused to other coding or regulatory sequences and still be considered isolated.

For example, recombinant DNA molecules contained in a vector are considered isolated. Further examples of isolated DNA molecules include recombinant DNA molecules maintained  
30 in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the isolated DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Accordingly, the present invention provides nucleic acid molecules that consist of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, cDNA sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists of a nucleotide sequence when the nucleotide  
5 sequence is the complete nucleotide sequence of the nucleic acid molecule.

The present invention further provides nucleic acid molecules that consist essentially of the nucleotide sequence shown in Figure 1 or 3 (SEQ ID NO:1, cDNA sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in Figure 2, SEQ ID NO:2. A nucleic acid molecule consists essentially of a nucleotide sequence  
10 when such a nucleotide sequence is present with only a few additional nucleic acid residues in the final nucleic acid molecule.

The present invention further provides nucleic acid molecules that comprise the nucleotide sequences shown in Figure 1 or 3 (SEQ ID NO:1, cDNA sequence and SEQ ID NO:3, genomic sequence), or any nucleic acid molecule that encodes the protein provided in  
15 Figure 2, SEQ ID NO:2. A nucleic acid molecule comprises a nucleotide sequence when the nucleotide sequence is at least part of the final nucleotide sequence of the nucleic acid molecule. In such a fashion, the nucleic acid molecule can be only the nucleotide sequence or have additional nucleic acid residues, such as nucleic acid residues that are naturally associated with it or heterologous nucleotide sequences. Such a nucleic acid molecule can have a few additional  
20 nucleotides or can comprises several hundred or more additional nucleotides. A brief description of how various types of these nucleic acid molecules can be readily made/isolated is provided below.

In Figures 1 and 3, both coding and non-coding sequences are provided. Because of the source of the present invention, human genomic sequences (Figure 3) and  
25 cDNA/transcript sequences (Figure 1), the nucleic acid molecules in the Figures will contain genomic intronic sequences, 5' and 3' non-coding sequences, gene regulatory regions and non-coding intergenic sequences. In general such sequence features are either noted in Figures 1 and 3 or can readily be identified using computational tools known in the art. As discussed below, some of the non-coding regions, particularly gene regulatory elements such  
30 as promoters, are useful for a variety of purposes, e.g. control of heterologous gene expression, target for identifying gene activity modulating compounds, and are particularly claimed as fragments of the genomic sequence provided herein.

The isolated nucleic acid molecules can encode the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature peptide (when the

mature form has more than one peptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, facilitate protein trafficking, prolong or shorten protein half-life or facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in situ*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

As mentioned above, the isolated nucleic acid molecules include, but are not limited to, the sequence encoding the GPCR peptide alone, the sequence encoding the mature peptide and additional coding sequences, such as a leader or secretory sequence (e.g., a pre-pro or pro-protein sequence), the sequence encoding the mature peptide, with or without the additional coding sequences, plus additional non-coding sequences, for example introns and non-coding 5' and 3' sequences such as transcribed but non-translated sequences that play a role in transcription, mRNA processing (including splicing and polyadenylation signals), ribosome binding and stability of mRNA. In addition, the nucleic acid molecule may be fused to a marker sequence encoding, for example, a peptide that facilitates purification.

Isolated nucleic acid molecules can be in the form of RNA, such as mRNA, or in the form DNA, including cDNA and genomic DNA obtained by cloning or produced by chemical synthetic techniques or by a combination thereof. The nucleic acid, especially DNA, can be double-stranded or single-stranded. Single-stranded nucleic acid can be the coding strand (sense strand) or the non-coding strand (anti-sense strand).

The invention further provides nucleic acid molecules that encode fragments of the peptides of the present invention as well as nucleic acid molecules that encode obvious variants of the GPCR proteins of the present invention that are described above. Such nucleic acid molecules may be naturally occurring, such as allelic variants (same locus), paralogs (different locus), and orthologs (different organism), or may be constructed by recombinant DNA methods or by chemical synthesis. Such non-naturally occurring variants may be made by mutagenesis techniques, including those applied to nucleic acid molecules, cells, or organisms. Accordingly, as discussed above, the variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions.

The present invention further provides non-coding fragments of the nucleic acid molecules provided in Figures 1 and 3. Preferred non-coding fragments include, but are not limited to, promoter sequences, enhancer sequences, gene modulating sequences and gene termination sequences. Such fragments are useful in controlling heterologous gene expression

and in developing screens to identify gene-modulating agents. A promoter can readily be identified as being 5' to the ATG start site in the genomic sequence provided in Figure 3.

A fragment comprises a contiguous nucleotide sequence greater than 12 or more nucleotides. Further, a fragment could be at least 30, 40, 50, 100, 250 or 500 nucleotides in length.

5 The length of the fragment will be based on its intended use. For example, the fragment can encode epitope bearing regions of the peptide, or can be useful as DNA probes and primers. Such fragments can be isolated using the known nucleotide sequence to synthesize an oligonucleotide probe. A labeled probe can then be used to screen a cDNA library, genomic DNA library, or mRNA to isolate nucleic acid corresponding to the coding region. Further,  
10 primers can be used in PCR reactions to clone specific regions of gene.

A probe/primer typically comprises substantially a purified oligonucleotide or oligonucleotide pair. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 20, 25, 40, 50 or more consecutive nucleotides.

15 Orthologs, homologs, and allelic variants can be identified using methods well known in the art. As described in the Peptide Section, these variants comprise a nucleotide sequence encoding a peptide that is typically 60-70%, 70-80%, 80-90%, and more typically at least about 90-95% or more homologous to the nucleotide sequence shown in the Figure sheets or a fragment of this sequence. Such nucleic acid molecules can readily be identified as being able  
20 to hybridize under moderate to stringent conditions, to the nucleotide sequence shown in the Figure sheets or a fragment of the sequence. Allelic variants can readily be determined by genetic locus of the encoding gene. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 1.

Figure 3 provides information on SNPs that have been found in a gene encoding the  
25 GPCR proteins of the present invention. The following SNPs were found: T406C, T852C, G897A, C1433T, T5845C, and G7028A.

As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences encoding a peptide at least 60-70% homologous to each other typically remain hybridized to each other. The  
30 conditions can be such that sequences at least about 60%, at least about 70%, or at least about 80% or more homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. One example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about

45C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65C. Examples of moderate to low stringency hybridization conditions are well known in the art.

#### Nucleic Acid Molecule Uses

5       The nucleic acid molecules of the present invention are useful for probes, primers, chemical intermediates, and in biological assays. The nucleic acid molecules are useful as a hybridization probe for messenger RNA, transcript/cDNA and genomic DNA to isolate full-length cDNA and genomic clones encoding the peptide described in Figure 2 and to isolate cDNA and genomic clones that correspond to variants (alleles, orthologs, etc.) producing the  
10       same or related peptides shown in Figure 2. As illustrated in Figure 3, known SNP variations include T406C, T852C, G897A, C1433T, T5845C, and G7028A.

      The probe can correspond to any sequence along the entire length of the nucleic acid molecules provided in the Figures. Accordingly, it could be derived from 5' noncoding regions, the coding region, and 3' noncoding regions. However, as discussed, fragments are not to be  
15       construed as encompassing fragments disclosed prior to the present invention.

      The nucleic acid molecules are also useful as primers for PCR to amplify any given region of a nucleic acid molecule and are useful to synthesize antisense molecules of desired length and sequence.

      The nucleic acid molecules are also useful for constructing recombinant vectors. Such  
20       vectors include expression vectors that express a portion of, or all of, the peptide sequences. Vectors also include insertion vectors, used to integrate into another nucleic acid molecule sequence, such as into the cellular genome, to alter *in situ* expression of a gene and/or gene product. For example, an endogenous coding sequence can be replaced via homologous recombination with all or part of the coding region containing one or more specifically  
25       introduced mutations.

      The nucleic acid molecules are also useful for expressing antigenic portions of the proteins.

      The nucleic acid molecules are also useful as probes for determining the chromosomal positions of the nucleic acid molecules by means of *in situ* hybridization methods. As indicated  
30       by the data presented in Figure 3, the map position was determined to be on chromosome 1.

      The nucleic acid molecules are also useful in making vectors containing the gene regulatory regions of the nucleic acid molecules of the present invention.

The nucleic acid molecules are also useful for designing ribozymes corresponding to all, or a part, of the mRNA produced from the nucleic acid molecules described herein.

The nucleic acid molecules are also useful for making vectors that express part, or all, of the peptides.

5       The nucleic acid molecules are also useful for constructing host cells expressing a part, or all, of the nucleic acid molecules and peptides.

The nucleic acid molecules are also useful for constructing transgenic animals expressing all, or a part, of the nucleic acid molecules and peptides.

10       The nucleic acid molecules are also useful as hybridization probes for determining the presence, level, form and distribution of nucleic acid expression. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Specifically, a virtual northern blot shows expression in a pooled fetal lung/testis/B-cell sample. In addition, PCR-based tissue screening panels indicate expression in humans in Hela cells and bone marrow.

15       Accordingly, the probes can be used to detect the presence of, or to determine levels of, a specific nucleic acid molecule in cells, tissues, and in organisms. The nucleic acid whose level is determined can be DNA or RNA. Accordingly, probes corresponding to the peptides described herein can be used to assess expression and/or gene copy number in a given cell, tissue, or organism. These uses are relevant for diagnosis of disorders involving an increase or

20       decrease in GPCR protein expression relative to normal results.

*In vitro* techniques for detection of mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detecting DNA includes Southern hybridizations and *in situ* hybridization.

25       Probes can be used as a part of a diagnostic test kit for identifying cells or tissues that express a GPCR protein, such as by measuring a level of a receptor-encoding nucleic acid in a sample of cells from a subject e.g., mRNA or genomic DNA, or determining if a receptor gene has been mutated. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Specifically, a virtual northern blot shows expression in a pooled fetal

30       lung/testis/B-cell sample. In addition, PCR-based tissue screening panels indicate expression in humans in Hela cells and bone marrow.

Nucleic acid expression assays are useful for drug screening to identify compounds that modulate GPCR nucleic acid expression.



The invention thus provides a method for identifying a compound that can be used to treat a disorder associated with nucleic acid expression of the GPCR gene, particularly biological and pathological processes that are mediated by the GPCR in cells and tissues that express it. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. The method typically includes assaying the ability of the compound to modulate the expression of the GPCR nucleic acid and thus identifying a compound that can be used to treat a disorder characterized by undesired GPCR nucleic acid expression. The assays can be performed in cell-based and cell-free systems. Cell-based assays include cells naturally expressing the GPCR nucleic acid or recombinant cells genetically engineered to express specific nucleic acid sequences.

The assay for GPCR nucleic acid expression can involve direct assay of nucleic acid levels, such as mRNA levels, or on collateral compounds involved in the signal pathway. Further, the expression of genes that are up- or down-regulated in response to the GPCR protein signal pathway can also be assayed. In this embodiment the regulatory regions of these genes can be operably linked to a reporter gene such as luciferase.

Thus, modulators of GPCR gene expression can be identified in a method wherein a cell is contacted with a candidate compound and the expression of mRNA determined. The level of expression of GPCR mRNA in the presence of the candidate compound is compared to the level of expression of GPCR mRNA in the absence of the candidate compound. The candidate compound can then be identified as a modulator of nucleic acid expression based on this comparison and be used, for example to treat a disorder characterized by aberrant nucleic acid expression. When expression of mRNA is statistically significantly greater in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of nucleic acid expression. When nucleic acid expression is statistically significantly less in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of nucleic acid expression.

The invention further provides methods of treatment, with the nucleic acid as a target, using a compound identified through drug screening as a gene modulator to modulate GPCR nucleic acid expression, particularly to modulate activities within a cell or tissue that expresses the proteins. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Specifically, a virtual northern blot shows expression in a pooled fetal lung/testis/B-cell sample. In addition, PCR-based tissue screening panels indicate expression in

humans in Hela cells and bone marrow. Modulation includes both up-regulation (i.e. activation or agonization) or down-regulation (suppression or antagonization) or nucleic acid expression.

Alternatively, a modulator for GPCR nucleic acid expression can be a small molecule or drug identified using the screening assays described herein as long as the drug or small molecule inhibits the GPCR nucleic acid expression in the cells and tissues that express the protein. Experimental data as provided in Figure 1 indicates expression in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells.

The nucleic acid molecules are also useful for monitoring the effectiveness of modulating compounds on the expression or activity of the GPCR gene in clinical trials or in a treatment regimen. Thus, the gene expression pattern can serve as a barometer for the continuing effectiveness of treatment with the compound, particularly with compounds to which a patient can develop resistance. The gene expression pattern can also serve as a marker indicative of a physiological response of the affected cells to the compound. Accordingly, such monitoring would allow either increased administration of the compound or the administration of alternative compounds to which the patient has not become resistant. Similarly, if the level of nucleic acid expression falls below a desirable level, administration of the compound could be commensurately decreased.

The nucleic acid molecules are also useful in diagnostic assays for qualitative changes in GPCR nucleic acid, and particularly in qualitative changes that lead to pathology. The nucleic acid molecules can be used to detect mutations in GPCR genes and gene expression products such as mRNA. The nucleic acid molecules can be used as hybridization probes to detect naturally-occurring genetic mutations in the GPCR gene and thereby to determine whether a subject with the mutation is at risk for a disorder caused by the mutation. Mutations include deletion, addition, or substitution of one or more nucleotides in the gene, chromosomal rearrangement, such as inversion or transposition, modification of genomic DNA, such as aberrant methylation patterns or changes in gene copy number, such as amplification. Detection of a mutated form of the GPCR gene associated with a dysfunction provides a diagnostic tool for an active disease or susceptibility to disease when the disease results from overexpression, underexpression, or altered expression of a GPCR protein.

Individuals carrying mutations in the GPCR gene can be detected at the nucleic acid level by a variety of techniques. Figure 3 provides information on SNPs that have been found in a gene encoding the GPCR proteins of the present invention. The following SNPs were found: T406C, T852C, G897A, C1433T, T5845C, and G7028A. As indicated by the data presented in Figure 3, the map position was determined to be on chromosome 1. Genomic DNA can be

analyzed directly or can be amplified by using PCR prior to analysis. RNA or cDNA can be used in the same way. In some uses, detection of the mutation involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.*, *Science* 241:1077-1080 (1988); and Nakazawa *et al.*, *PNAS* 91:360-364 (1994)), the latter of which can be particularly useful for detecting point mutations in the gene (see Abravaya *et al.*, *Nucleic Acids Res.* 23:675-682 (1995)). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a gene under conditions such that hybridization and amplification of the gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. Deletions and insertions can be detected by a change in size of the amplified product compared to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to normal RNA or antisense DNA sequences.

Alternatively, mutations in a GPCR gene can be directly identified, for example, by alterations in restriction enzyme digestion patterns determined by gel electrophoresis.

Further, sequence-specific ribozymes (U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

Perfectly matched sequences can be distinguished from mismatched sequences by nuclease cleavage digestion assays or by differences in melting temperature.

Sequence changes at specific locations can also be assessed by nuclease protection assays such as RNase and S1 protection or the chemical cleavage method. Furthermore, sequence differences between a mutant GPCR gene and a wild-type gene can be determined by direct DNA sequencing. A variety of automated sequencing procedures can be utilized when performing the diagnostic assays (Naeve, C.W., (1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.*, *Adv. Chromatogr.* 36:127-162 (1996); and Griffin *et al.*, *Appl. Biochem. Biotechnol.* 38:147-159 (1993)).

Other methods for detecting mutations in the gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA duplexes (Myers *et al.*, *Science* 230:1242 (1985)); Cotton *et al.*, *PNAS* 85:4397 (1988); Saleeba *et al.*, *Meth. Enzymol.* 217:286-295 (1992)), electrophoretic mobility of mutant and wild type nucleic acid is compared (Orita *et al.*, *PNAS* 86:2766 (1989); Cotton *et al.*, *Mutat. Res.* 285:125-144

(1993); and Hayashi *et al.*, *Genet. Anal. Tech. Appl.* 9:73-79 (1992)), and movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (Myers *et al.*, *Nature* 313:495 (1985)). Examples of other techniques for detecting point mutations include selective oligonucleotide hybridization, selective amplification, and selective primer extension.

The nucleic acid molecules are also useful for testing an individual for a genotype that while not necessarily causing the disease, nevertheless affects the treatment modality. Thus, the nucleic acid molecules can be used to study the relationship between an individual's genotype and the individual's response to a compound used for treatment (pharmacogenomic relationship). Accordingly, the nucleic acid molecules described herein can be used to assess the mutation content of the GPCR gene in an individual in order to select an appropriate compound or dosage regimen for treatment. As illustrated in Figure 3, known SNP variations include T406C, T852C, G897A, C1433T, T5845C, and G7028A.

Thus nucleic acid molecules displaying genetic variations that affect treatment provide a diagnostic target that can be used to tailor treatment in an individual. Accordingly, the production of recombinant cells and animals containing these polymorphisms allow effective clinical design of treatment compounds and dosage regimens.

The nucleic acid molecules are thus useful as antisense constructs to control GPCR gene expression in cells, tissues, and organisms. A DNA antisense nucleic acid molecule is designed to be complementary to a region of the gene involved in transcription, preventing transcription and hence production of GPCR protein. An antisense RNA or DNA nucleic acid molecule would hybridize to the mRNA and thus block translation of mRNA into GPCR protein.

Alternatively, a class of antisense molecules can be used to inactivate mRNA in order to decrease expression of GPCR nucleic acid. Accordingly, these molecules can treat a disorder characterized by abnormal or undesired GPCR nucleic acid expression. This technique involves cleavage by means of ribozymes containing nucleotide sequences complementary to one or more regions in the mRNA that attenuate the ability of the mRNA to be translated. Possible regions include coding regions and particularly coding regions corresponding to the catalytic and other functional activities of the GPCR protein, such as ligand binding.

The nucleic acid molecules also provide vectors for gene therapy in patients containing cells that are aberrant in GPCR gene expression. Thus, recombinant cells, which include the patient's cells that have been engineered *ex vivo* and returned to the patient, are introduced into an individual where the cells produce the desired GPCR protein to treat the individual.

The invention also encompasses kits for detecting the presence of a GPCR nucleic acid in a biological sample. Experimental data as provided in Figure 1 indicates that GPCR proteins of the present invention are expressed in Hela cells, bone marrow, and a pooled sample of fetal lung, testis, and B-cells. Specifically, a virtual northern blot shows expression in a pooled fetal lung/testis/B-cell sample. In addition, PCR-based tissue screening panels indicate expression in humans in Hela cells and bone marrow. For example, the kit can comprise reagents such as a labeled or labelable nucleic acid or agent capable of detecting GPCR nucleic acid in a biological sample; means for determining the amount of GPCR nucleic acid in the sample; and means for comparing the amount of GPCR nucleic acid in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect GPCR protein mRNA or DNA.

#### Nucleic Acid Arrays

The present invention further provides nucleic acid detection kits, such as arrays or microarrays of nucleic acid molecules that are based on the sequence information provided in Figures 1 and 3 (SEQ ID NOS:1 and 3).

As used herein "Arrays" or "Microarrays" refers to an array of distinct polynucleotides or oligonucleotides synthesized on a substrate, such as paper, nylon or other type of membrane, filter, chip, glass slide, or any other suitable solid support. In one embodiment, the microarray is prepared and used according to the methods described in US Patent 5,837,832, Chee et al., PCT application W095/11995 (Chee et al.), Lockhart, D. J. et al. (1996; Nat. Biotech. 14: 1675-1680) and Schena, M. et al. (1996; Proc. Natl. Acad. Sci. 93: 10614-10619), all of which are incorporated herein in their entirety by reference. In other embodiments, such arrays are produced by the methods described by Brown et. al., US Patent No. 5,807,522.

The microarray or detection kit is preferably composed of a large number of unique, single-stranded nucleic acid sequences, usually either synthetic antisense oligonucleotides or fragments of cDNAs, fixed to a solid support. The oligonucleotides are preferably about 6-60 nucleotides in length, more preferably 15-30 nucleotides in length, and most preferably about 20-25 nucleotides in length. For a certain type of microarray or detection kit, it may be preferable to use oligonucleotides that are only 7-20 nucleotides in length. The microarray or detection kit may contain oligonucleotides that cover the known 5', or 3', sequence, sequential oligonucleotides which cover the full length sequence; or unique oligonucleotides selected

from particular areas along the length of the sequence. Polynucleotides used in the microarray or detection kit may be oligonucleotides that are specific to a gene or genes of interest.

In order to produce oligonucleotides to a known sequence for a microarray or detection kit, the gene(s) of interest (or an ORF identified from the contigs of the present invention) is typically examined using a computer algorithm which starts at the 5' or at the 3' end of the nucleotide sequence. Typical algorithms will then identify oligomers of defined length that are unique to the gene, have a GC content within a range suitable for hybridization, and lack predicted secondary structure that may interfere with hybridization. In certain situations it may be appropriate to use pairs of oligonucleotides on a microarray or detection kit. The "pairs" will be identical, except for one nucleotide that preferably is located in the center of the sequence. The second oligonucleotide in the pair (mismatched by one) serves as a control. The number of oligonucleotide pairs may range from two to one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support.

In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet application apparatus, as described in PCT application W095/251116 (Baldeschweiler et al.) which is incorporated herein in its entirety by reference. In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536, 6144 or more oligonucleotides, or any other number between two and one million which lends itself to the efficient use of commercially available instrumentation.

In order to conduct sample analysis using a microarray or detection kit, the RNA or DNA from a biological sample is made into hybridization probes. The mRNA is isolated, and cDNA is produced and used as a template to make antisense RNA (aRNA). The aRNA is amplified in the presence of fluorescent nucleotides, and labeled probes are incubated with the microarray or detection kit so that the probe sequences hybridize to complementary oligonucleotides of the microarray or detection kit. Incubation conditions are adjusted so that hybridization occurs with precise complementary matches or with various degrees of less complementarity. After removal of nonhybridized probes, a scanner is used to determine the

levels and patterns of fluorescence. The scanned images are examined to determine degree of complementarity and the relative abundance of each oligonucleotide sequence on the microarray or detection kit. The biological samples may be obtained from any bodily fluids (such as blood, urine, saliva, phlegm, gastric juices, etc.), cultured cells, biopsies, or other  
5 tissue preparations. A detection system may be used to measure the absence, presence, and amount of hybridization for all of the distinct sequences simultaneously. This data may be used for large scale correlation studies on the sequences, expression patterns, mutations, variants, or polymorphisms among samples.

Using such arrays, the present invention provides methods to identify the expression  
10 of the GPCR proteins/peptides of the present invention. In detail, such methods comprise incubating a test sample with one or more nucleic acid molecules and assaying for binding of the nucleic acid molecule with components within the test sample. Such assays will typically involve arrays comprising many genes, at least one of which is a gene of the present invention and or alleles of the GPCR gene of the present invention. Figure 3 provides  
15 information on SNPs that have been found in a gene encoding the GPCR proteins of the present invention. The following SNPs were found: T406C, T852C, G897A, C1433T, T5845C, and G7028A.

Conditions for incubating a nucleic acid molecule with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and  
20 the type and nature of the nucleic acid molecule used in the assay. One skilled in the art will recognize that any one of the commonly available hybridization, amplification or array assay formats can readily be adapted to employ the novel fragments of the Human genome disclosed herein. Examples of such assays can be found in Chard, T, *An Introduction to Radioimmunoassay and Related Techniques*, Elsevier Science Publishers, Amsterdam, The  
25 Netherlands (1986); Bullock, G. R. *et al.*, *Techniques in Immunocytochemistry*, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., *Practice and Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology*, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The test samples of the present invention include cells, protein or membrane extracts  
30 of cells. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing nucleic acid extracts or of cells are well known in the art and can be readily be adapted in order to obtain a sample that is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention.

Specifically, the invention provides a compartmentalized kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the nucleic acid molecules that can bind to a fragment of the Human genome disclosed herein; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound nucleic acid.

In detail, a compartmentalized kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers, strips of plastic, glass or paper, or arraying material such as silica. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the nucleic acid probe, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound probe. One skilled in the art will readily recognize that the previously unidentified GPCR genes of the present invention can be routinely identified using the sequence information disclosed herein can be readily incorporated into one of the established kit formats which are well known in the art, particularly expression arrays.

#### Vectors/host cells

The invention also provides vectors containing the nucleic acid molecules described herein. The term "vector" refers to a vehicle, preferably a nucleic acid molecule, which can transport the nucleic acid molecules. When the vector is a nucleic acid molecule, the nucleic acid molecules are covalently linked to the vector nucleic acid. With this aspect of the invention, the vector includes a plasmid, single or double stranded phage, a single or double stranded RNA or DNA viral vector, or artificial chromosome, such as a BAC, PAC, YAC, OR MAC.

A vector can be maintained in the host cell as an extrachromosomal element where it replicates and produces additional copies of the nucleic acid molecules. Alternatively, the vector may integrate into the host cell genome and produce additional copies of the nucleic acid molecules when the host cell replicates.



The invention provides vectors for the maintenance (cloning vectors) or vectors for expression (expression vectors) of the nucleic acid molecules. The vectors can function in procaryotic or eukaryotic cells or in both (shuttle vectors).

Expression vectors contain cis-acting regulatory regions that are operably linked in the vector to the nucleic acid molecules such that transcription of the nucleic acid molecules is allowed in a host cell. The nucleic acid molecules can be introduced into the host cell with a separate nucleic acid molecule capable of affecting transcription. Thus, the second nucleic acid molecule may provide a trans-acting factor interacting with the cis-regulatory control region to allow transcription of the nucleic acid molecules from the vector. Alternatively, a trans-acting factor may be supplied by the host cell. Finally, a trans-acting factor can be produced from the vector itself. It is understood, however, that in some embodiments, transcription and/or translation of the nucleic acid molecules can occur in a cell-free system.

The regulatory sequence to which the nucleic acid molecules described herein can be operably linked include promoters for directing mRNA transcription. These include, but are not limited to, the left promoter from bacteriophage  $\lambda$ , the lac, TRP, and TAC promoters from *E. coli*, the early and late promoters from SV40, the CMV immediate early promoter, the adenovirus early and late promoters, and retrovirus long-terminal repeats.

In addition to control regions that promote transcription, expression vectors may also include regions that modulate transcription, such as repressor binding sites and enhancers.

Examples include the SV40 enhancer, the cytomegalovirus immediate early enhancer, polyoma enhancer, adenovirus enhancers, and retrovirus LTR enhancers.

In addition to containing sites for transcription initiation and control, expression vectors can also contain sequences necessary for transcription termination and, in the transcribed region a ribosome binding site for translation. Other regulatory control elements for expression include initiation and termination codons as well as polyadenylation signals. The person of ordinary skill in the art would be aware of the numerous regulatory sequences that are useful in expression vectors. Such regulatory sequences are described, for example, in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

A variety of expression vectors can be used to express a nucleic acid molecule. Such vectors include chromosomal, episomal, and virus-derived vectors, for example vectors derived from bacterial plasmids, from bacteriophage, from yeast episomes, from yeast chromosomal elements, including yeast artificial chromosomes, from viruses such as baculoviruses,

papovaviruses such as SV40, Vaccinia viruses, adenoviruses, poxviruses, pseudorabies viruses, and retroviruses. Vectors may also be derived from combinations of these sources such as those derived from plasmid and bacteriophage genetic elements, eg. cosmids and phagemids.

Appropriate cloning and expression vectors for prokaryotic and eukaryotic hosts are described in  
5 Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*. 2nd. ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1989).

The regulatory sequence may provide constitutive expression in one or more host cells (i.e. tissue specific) or may provide for inducible expression in one or more cell types such as by temperature, nutrient additive, or exogenous factor such as a hormone or other ligand. A variety  
10 of vectors providing for constitutive and inducible expression in prokaryotic and eukaryotic hosts are well known to those of ordinary skill in the art.

The nucleic acid molecules can be inserted into the vector nucleic acid by well-known methodology. Generally, the DNA sequence that will ultimately be expressed is joined to an expression vector by cleaving the DNA sequence and the expression vector with one or more  
15 restriction enzymes and then ligating the fragments together. Procedures for restriction enzyme digestion and ligation are well known to those of ordinary skill in the art.

The vector containing the appropriate nucleic acid molecule can be introduced into an appropriate host cell for propagation or expression using well-known techniques. Bacterial cells include, but are not limited to, *E. coli*, *Streptomyces*, and *Salmonella typhimurium*. Eukaryotic  
20 cells include, but are not limited to, yeast, insect cells such as *Drosophila*, animal cells such as COS and CHO cells, and plant cells.

As described herein, it may be desirable to express the peptide as a fusion protein. Accordingly, the invention provides fusion vectors that allow for the production of the peptides. Fusion vectors can increase the expression of a recombinant protein, increase the solubility of  
25 the recombinant protein, and aid in the purification of the protein by acting for example as a ligand for affinity purification. A proteolytic cleavage site may be introduced at the junction of the fusion moiety so that the desired peptide can ultimately be separated from the fusion moiety. Proteolytic enzymes include, but are not limited to, factor Xa, thrombin, and enterokinase. Typical fusion expression vectors include pGEX (Smith *et al.*, *Gene* 67:31-40 (1988)), pMAL  
30 (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein. Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, *Gene* 69:301-315 (1988)) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185:60-89 (1990)).

Recombinant protein expression can be maximized in a host bacteria by providing a genetic background wherein the host cell has an impaired capacity to proteolytically cleave the recombinant protein. (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Alternatively, the sequence of the nucleic acid molecule of interest can be altered to provide preferential codon usage for a specific host cell, for example *E. coli*. (Wada *et al.*, *Nucleic Acids Res.* 20:2111-2118 (1992)).

The nucleic acid molecules can also be expressed by expression vectors that are operative in yeast. Examples of vectors for expression in yeast e.g., *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, *EMBO J.* 6:229-234 (1987)), pMFa (Kurjan *et al.*, *Cell* 30:933-943(1982)), pJRY88 (Schultz *et al.*, *Gene* 54:113-123 (1987)), and pYES2 (Invitrogen Corporation, San Diego, CA).

The nucleic acid molecules can also be expressed in insect cells using, for example, baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.*, *Mol. Cell Biol.* 3:2156-2165 (1983)) and the pVL series (Lucklow *et al.*, *Virology* 170:31-39 (1989)).

In certain embodiments of the invention, the nucleic acid molecules described herein are expressed in mammalian cells using mammalian expression vectors. Examples of mammalian expression vectors include pCDM8 (Seed, B. *Nature* 329:840(1987)) and pMT2PC (Kaufman *et al.*, *EMBO J.* 6:187-195 (1987)).

The expression vectors listed herein are provided by way of example only of the well-known vectors available to those of ordinary skill in the art that would be useful to express the nucleic acid molecules. The person of ordinary skill in the art would be aware of other vectors suitable for maintenance propagation or expression of the nucleic acid molecules described herein. These are found for example in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed.*, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

The invention also encompasses vectors in which the nucleic acid sequences described herein are cloned into the vector in reverse orientation, but operably linked to a regulatory sequence that permits transcription of antisense RNA. Thus, an antisense transcript can be produced to all, or to a portion, of the nucleic acid molecule sequences described herein, including both coding and non-coding regions. Expression of this antisense RNA is subject to each of the parameters described above in relation to expression of the sense RNA (regulatory sequences, constitutive or inducible expression, tissue-specific expression).

The invention also relates to recombinant host cells containing the vectors described herein. Host cells therefore include prokaryotic cells, lower eukaryotic cells such as yeast, other eukaryotic cells such as insect cells, and higher eukaryotic cells such as mammalian cells.

The recombinant host cells are prepared by introducing the vector constructs described  
5 herein into the cells by techniques readily available to the person of ordinary skill in the art. These include, but are not limited to, calcium phosphate transfection, DEAE-dextran-mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, lipofection, and other techniques such as those found in Sambrook, *et al.* (*Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory  
10 Press, Cold Spring Harbor, NY, 1989).

Host cells can contain more than one vector. Thus, different nucleotide sequences can be introduced on different vectors of the same cell. Similarly, the nucleic acid molecules can be introduced either alone or with other nucleic acid molecules that are not related to the nucleic acid molecules such as those providing trans-acting factors for expression vectors. When more  
15 than one vector is introduced into a cell, the vectors can be introduced independently, co-introduced or joined to the nucleic acid molecule vector.

In the case of bacteriophage and viral vectors, these can be introduced into cells as packaged or encapsulated virus by standard procedures for infection and transduction. Viral vectors can be replication-competent or replication-defective. In the case in which viral  
20 replication is defective, replication will occur in host cells providing functions that complement the defects.

Vectors generally include selectable markers that enable the selection of the subpopulation of cells that contain the recombinant vector constructs. The marker can be contained in the same vector that contains the nucleic acid molecules described herein or may be  
25 on a separate vector. Markers include tetracycline or ampicillin-resistance genes for prokaryotic host cells and dihydrofolate reductase or neomycin resistance for eukaryotic host cells. However, any marker that provides selection for a phenotypic trait will be effective.

While the mature proteins can be produced in bacteria, yeast, mammalian cells, and other cells under the control of the appropriate regulatory sequences, cell-free transcription and  
30 translation systems can also be used to produce these proteins using RNA derived from the DNA constructs described herein.

Where secretion of the peptide is desired, which is difficult to achieve with multi-transmembrane domain containing proteins such as GPCRs, appropriate secretion signals are

incorporated into the vector. The signal sequence can be endogenous to the peptides or heterologous to these peptides.

Where the peptide is not secreted into the medium, which is typically the case with GPCRs, the protein can be isolated from the host cell by standard disruption procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. The peptide can then be recovered and purified by well-known purification methods including ammonium sulfate precipitation, acid extraction, anion or cationic exchange chromatography, phosphocellulose chromatography, hydrophobic-interaction chromatography, affinity chromatography, hydroxylapatite chromatography, lectin chromatography, or high performance liquid chromatography.

It is also understood that depending upon the host cell in recombinant production of the peptides described herein, the peptides can have various glycosylation patterns, depending upon the cell, or maybe non-glycosylated as when produced in bacteria. In addition, the peptides may include an initial modified methionine in some cases as a result of a host-mediated process.

#### Uses of vectors and host cells

The recombinant host cells expressing the peptides described herein have a variety of uses. First, the cells are useful for producing a GPCR protein or peptide that can be further purified to produce desired amounts of GPCR protein or fragments. Thus, host cells containing expression vectors are useful for peptide production.

Host cells are also useful for conducting cell-based assays involving the GPCR protein or GPCR protein fragments, such as those described above as well as other formats known in the art. Thus, a recombinant host cell expressing a native GPCR protein is useful for assaying compounds that stimulate or inhibit GPCR protein function.

Host cells are also useful for identifying GPCR protein mutants in which these functions are affected. If the mutants naturally occur and give rise to a pathology, host cells containing the mutations are useful to assay compounds that have a desired effect on the mutant GPCR protein (for example, stimulating or inhibiting function) which may not be indicated by their effect on the native GPCR protein.

Genetically engineered host cells can be further used to produce non-human transgenic animals. A transgenic animal is preferably a mammal, for example a rodent, such as a rat or mouse, in which one or more of the cells of the animal include a transgene. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal

develops and which remains in the genome of the mature animal in one or more cell types or tissues of the transgenic animal. These animals are useful for studying the function of a GPCR protein and identifying and evaluating modulators of GPCR protein activity. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, and  
5 amphibians.

A transgenic animal can be produced by introducing nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. Any of the GPCR protein nucleotide sequences can be introduced as a transgene into the genome of a non-human animal, such as a  
10 mouse.

Any of the regulatory or other sequences useful in expression vectors can form part of the transgenic sequence. This includes intronic sequences and polyadenylation signals, if not already included. A tissue-specific regulatory sequence(s) can be operably linked to the transgene to direct expression of the GPCR protein to particular cells.

15 Methods for generating transgenic animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for  
20 production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the transgene in its genome and/or expression of transgenic mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene can further be bred to other transgenic animals carrying other transgenes. A transgenic animal also includes  
25 animals in which the entire animal or tissues in the animal have been produced using the homologously recombinant host cells described herein.

In another embodiment, transgenic non-human animals can be produced which contain selected systems that allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the  
30 *cre/loxP* recombinase system, see, e.g., Lakso *et al.* *PNAS* 89:6232-6236 (1992). Another example of a recombinase system is the FLP recombinase system of *S. cerevisiae* (O'Gorman *et al.* *Science* 251:1351-1355 (1991). If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein is required. Such animals can be provided through the construction of

"double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al. Nature* 385:810-813 (1997) and PCT International Publication Nos. WO 97/07668 and WO 97/07669. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G<sub>0</sub> phase. The quiescent cell can then be fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to pseudopregnant female foster animal. The offspring born of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

Transgenic animals containing recombinant cells that express the peptides described herein are useful to conduct the assays described herein in an *in vivo* context. Accordingly, the various physiological factors that are present *in vivo* and that could effect ligand binding, GPCR protein activation, and signal transduction, may not be evident from *in vitro* cell-free or cell-based assays. Accordingly, it is useful to provide non-human transgenic animals to assay *in vivo* GPCR protein function, including ligand interaction, the effect of specific mutant GPCR proteins on GPCR protein function and ligand interaction, and the effect of chimeric GPCR proteins. It is also possible to assess the effect of null mutations, that is mutations that substantially or completely eliminate one or more GPCR protein functions.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above-described modes for carrying out the invention which are obvious to those skilled in the field of molecular biology or related fields are intended to be within the scope of the following claims.

### Claims

That which is claimed is:

1. An isolated peptide consisting of an amino acid sequence selected from the group consisting of:
  - (a) an amino acid sequence shown in SEQ ID NO:2;
  - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
  - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic); and
  - (d) a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids.
2. An isolated peptide comprising an amino acid sequence selected from the group consisting of:
  - (a) an amino acid sequence shown in SEQ ID NO:2;
  - (b) an amino acid sequence of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said allelic variant is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
  - (c) an amino acid sequence of an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said ortholog is encoded by a nucleic acid molecule that hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic); and
  - (d) a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids.
3. An isolated antibody that selectively binds to a peptide of claim 2.



4. An isolated nucleic acid molecule consisting of a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NO:2;
- (b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
- (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
- (d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids; and
- (e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

5. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence that encodes an amino acid sequence shown in SEQ ID NO:2;
- (b) a nucleotide sequence that encodes of an allelic variant of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
- (c) a nucleotide sequence that encodes an ortholog of an amino acid sequence shown in SEQ ID NO:2, wherein said nucleotide sequence hybridizes under stringent conditions to the opposite strand of a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic);
- (d) a nucleotide sequence that encodes a fragment of an amino acid sequence shown in SEQ ID NO:2, wherein said fragment comprises at least 10 contiguous amino acids; and

(e) a nucleotide sequence that is the complement of a nucleotide sequence of (a)-(d).

6. A gene chip comprising a nucleic acid molecule of claim 5.
7. A transgenic non-human animal comprising a nucleic acid molecule of claim 5.
8. A nucleic acid vector comprising a nucleic acid molecule of claim 5.
9. A host cell containing the vector of claim 8.
10. A method for producing any of the peptides of claim 1 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
11. A method for producing any of the peptides of claim 2 comprising introducing a nucleotide sequence encoding any of the amino acid sequences in (a)-(d) into a host cell, and culturing the host cell under conditions in which the peptides are expressed from the nucleotide sequence.
12. A method for detecting the presence of any of the peptides of claim 2 in a sample, said method comprising contacting said sample with a detection agent that specifically allows detection of the presence of the peptide in the sample and then detecting the presence of the peptide.
13. A method for detecting the presence of a nucleic acid molecule of claim 5 in a sample, said method comprising contacting the sample with an oligonucleotide that hybridizes to said nucleic acid molecule under stringent conditions and determining whether the oligonucleotide binds to said nucleic acid molecule in the sample.
14. A method for identifying a modulator of a peptide of claim 2, said method comprising contacting said peptide with an agent and determining if said agent has modulated the function or activity of said peptide.

15. The method of claim 14, wherein said agent is administered to a host cell comprising an expression vector that expresses said peptide.

16. A method for identifying an agent that binds to any of the peptides of claim 2, said method comprising contacting the peptide with an agent and assaying the contacted mixture to determine whether a complex is formed with the agent bound to the peptide.

17. A pharmaceutical composition comprising an agent identified by the method of claim 16 and a pharmaceutically acceptable carrier therefor.

18. A method for treating a disease or condition mediated by a human proteases, said method comprising administering to a patient a pharmaceutically effective amount of an agent identified by the method of claim 16.

19. A method for identifying a modulator of the expression of a peptide of claim 2, said method comprising contacting a cell expressing said peptide with an agent, and determining if said agent has modulated the expression of said peptide.

20. An isolated human protease peptide having an amino acid sequence that shares at least 70% homology with an amino acid sequence shown in SEQ ID NO:2.

21. A peptide according to claim 20 that shares at least 90 percent homology with an amino acid sequence shown in SEQ ID NO:2.

22. An isolated nucleic acid molecule encoding a human protease peptide, said nucleic acid molecule sharing at least 80 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic).

23. A nucleic acid molecule according to claim 22 that shares at least 90 percent homology with a nucleic acid molecule shown in SEQ ID NOS:1 (transcript) or 3 (genomic).

1/10

```

1   CCCGGGCTGC AGGAATTCGG CACGAGGCCA TGCTGGGCCC TGCTGTCCTG
51  GGCCTCAGCC TCTGGGCTCT CCTGCACCCT GGGACGGGGG CCCCATTGTG
101 CCTGTCACAG CAACTTAGGA TGAAGGGGGA CTACGTGCTG GGGGGGCTGT
151 TCCCCCTGGG CGAGGCCGAG GAGGCTGGCC TCCGCAGCCG GACACGGCCC
201 AGCAGCCCTG TGTGCACCAG GTTCTCCTCA AACGGCCTGC TCTGGGCACT
251 GGCCATGAAA ATGGCCGTGG AGGAGATCAA CAACAAGTCG GATCTGCTGC
301 CCGGGCTGCG CCTGGGCTAC GACCTCTTTG ATACGTGCTC GGAGCCTGTG
351 GTGGCCATGA AGCCCAGCCT CATGTTCTTG GCCAAGGCAG GCAGCCGCGA
401 CATCGCCGCC TACTGCAACT ACACGCAGTA CCAGCCCCGT GTGCTGGCTG
451 TCATCGGGCC CCACTCGTCA GAGCTCGCCA TGGTCACCGG CAAGTTCTTC
501 AGCTTCTTCC TCATGCCCCA GGTGAGCTAC GGTGCTAGCA TGGAGCTGCT
551 GAGCGCCCGG GAGACCTTCC CCTCCTTCTT CCGCACCGTG CCCAGCGACC
601 GTGTGCAGCT GACGGCCGCC GCGGAGCTGC TGCAGGAGTT CGGCTGGAAC
651 TGGGTGGCCG CCCTGGGCAG CGACGACGAG TACGGCCGGC AGGGCCTGAG
701 CATCTTCTCG GCCCTGGCCG CGGCACGCGG CATCTGCATC GCGCACGAGG
751 GCCTGGTGCC GCTGCCCCGT GCCGATGACT CGCGGCTGGG GAAGGTGCAG
801 GACGTCTCTG ACCAGGTGAA CCAGAGCAGC GTGCAGGTGG TGCTGCTGTT
851 CGCCTCCGTG CACGCCGCCC ACGCCTCTT CAACTACAGC ATCAGCAGCA
901 GGCTCTCGCC CAAGGTGTGG GTGGCCAGCG AGGCCCTGGT GACCTCTGAC
951 CTGCTCATGG GGCTGCCCGG CATGGCCAG ATGGGCACGG TGCTTGGCTT
1001 CCTCCAGAGG GGTGCCCAGC TGCACGAGTT CCCCAGTAC GTGAAGACGC
1051 ACCTGGCCCT GGCCACCGAC CCGGCCTTCT GCTCTGCCCT GGGCGAGAGG
1101 GAGCAGGGTC TGGAGGAGGA CGTGGTGGGC CAGCGCTGCC CGCAGTGTGA
1151 CTGCATCAGC CTGCAGAACG TGAGCGCAGG GCTAAATCAC CACCAGACGT
1201 TCTCTGTCTA CGCAGCTGTG TATAGCGTGG CCCAGGCCCT GCACAACACT
1251 CTTCAGTGCA ACCCTCAGG CTGCCCCGCG CAGGACCCCG TGAAGCCCTG
1301 GCAGCTCCTG GAGAACATGT ACAACCTGAC CTTCACGTG GCGGGGCTGC
1351 CGCTGCGGTT CGACAGCAGC GGAACGTGG ACATGGAGTA CGACCTGAAG
1401 CTGTGGGTGT GGCAGGGCTC AGTGCCAGG CTCCACGACG TGGGCAGGTT
1451 CACCGCGAGC CTCAGGACAG AGCGCCTGAA GATCCGCTGG CACACGTCTG
1501 ACAACCAGAA GCCCGTGTCC CGGTGCTCGC GGCAGTGCCA GGAGGGCCAG
1551 GTGCGCCGGG TCAAGGGGTT CCACTCCTGC TGCTACGACT GTGTGGACTG
1601 CGAGGCGGGC AGCTACCGGC AAAACCCAGA CGACATCGCC TGCACCTTTT
1651 GTGGCCAGGA TGAGTGGTCC CCGGAGCGAA GCACACGCTG CTTCCGCCCG
1701 AGGTCTCGGT TCCTGGCATG GGGCGAGCGG GCTGTGCTGC TGCTGCTCCT
1751 GCTGCTGAGC CTGGCGCTGG GCCTTGCTGT GGTGCTTTG GGGCTGTTCTG
1801 TTCACCATCG GGACAGCCCA CTGGTTCAGG CCTCGGGGGG GCCCCTGGCC
1851 TGCTTTGGCC TGGTGTGCCT GGGCCTGGTC TGCCCTCAGCG TCCTCCTGTT
1901 CCCTGGCCAG CCCAGCCCTG CCCGATGCCT GGGCCAGCAG CCCTTGTCCTC
1951 ACCTCCCGCT CACGGGCTGC CTGAGCACAC TCTTCCTGCA GCGGGCCGAG
2001 ATCTTCGTGG AGTCAGAACT GCCTCTGAGC TGGGCAGACC GGCTGAGTGG
2051 CTGCCTGCGG GGGCCCTGGG CCTGGCTGGT GGTGCTGCTG GCCATGCTGG
2101 TGGAGGTCGC ACTGTGCACC TGGTACCTGG TGGCCTTCCC GCCCGAGGTG
2151 GTGACGGACT GGCACATGCT GCCACGGAG GCGCTGGTGC ACTGCCCGAC
2201 ACGCTCTTGG GTACGCTTCG GCCTAGCGCA GCCACCAAT GCCACGCTGG
2251 CCTTTCTCTG CTTCCTGGGC ACTTCCTGG TCGCGAGCCA GCCGGGCCGC
2301 TACAACCGTG CCCGTGGCCT CACCTTTGCC ATGCTGGCCT ACTTCATCAC
2351 CTGGGTCTCC TTTGTGCCCC TCCTGGCCAA TGTGCAGGTG GTCCTCAGGC
2401 CCGCCGTGCA GATGGGCGCC CTCCTGCTCT GTGTCTTGGG CATCCTGGCT
2451 GCCTTCCACC TGCCAGGTG TTACCTGCTC ATGCGGCAGC CAGGGCTCAA
2501 CACCCCGCAG TTCTTCTCTG GAGGGGGCCC TGGGGATGCC CAAGGCCAGA
2551 ATGACGGGAA CACAGGAAAT CAGGGGAAAC ATGAGTGACC CAACCCTGTG
2601 ATCTCAGCCC CGGTGAACCC AGACTTAGCT GCGATCCCCC CCAAGCCAGC
2651 AATGACCCGT GTCTCGCTAC AGAGACCCTC CCGCTCTAGG TTCTGACCCC
2701 AGGTTGTCTC CTGACCCTGA CCCCACAGTG AGCCCTAGGC CTGGAGCACG
2751 TGGACACCCC TGTGACCATC TGGGCCCCAG AGCCAAGCTG TGTCCCTGTC
2801 CCTCTGTGCC CAGACCAGGC CTGCCCAGGT AACCAGACC CACTGTTCTG
2851 GAAAGAGGCC CGGAGGGCTC CCAGGGTACC CGCAACCCAC ACCGTGAGCT
2901 CAGGAAAAGG ACGCAGGGAG GCCCCGGCCA GATGGCTGGA AGCCCAAATC
2951 AGGCCCTGCC GACCTGACCA TGTCCACCA GGGCCCCAT CCTGCACCTT
3001 GCCAGGCACC ACAGCAGTGG GAGGCCAGGT GGGGGCACAC AGGCATATGC
3051 CCAGGGCAGA GCCCCCGGAG GTGGGGGTGG CACCCAGCTT CCTACTCTGC
3101 CCTTGCCCCA GTGGGTAGAC AGCATCATGA CTGTCACCAG TACCAGGGAC
3151 AGAGCCCAGG TGGGGTGGGG GCGGGGTCCA GCACCACGGC CAGCACCAGC
3201 CACCAAGACC CCGGAGCCAG CACCATGGAC AGAAAACCTG CCACCAGGAT
3251 CTGACGCCAG CACGCCGCCA GGCCACACA GGGTCTCCGG TCAGAGTCCC
3301 AGGGTCAGCT CCCAGCAGGG CCTAGGGGAG GCTGGACCAG CTCCTGTGCTG

```

FIGURE 1, page 1 of 2

2/10

3351 CTCATTCCAA GGCAGCCCAG CCGGAGAGAA GGGGCACAGG CCACACATCT  
 3401 GTCCCATAAA ATTAAACGCT TTTTAGTGTT TAAAATAAAA AAAAAAAAAA  
 3451 AAAAAAAAAA  
 (SEQ ID NO:1)

**FEATURES:**

Start: 30  
 Stop: 2586

**HOMOLOGOUS PROTEIN:**Top 10 BLAST Hits:

	Score	E
gi 4337086 gb AAD18069.1  (AF127389) putative taste receptor TR...	460	e-128
gi 1168781 sp P41180 CASR_HUMAN EXTRACELLULAR CALCIUM-SENSING R...	395	e-109
gi 4557411 ref NP_000379.1  calcium-sensing receptor (hypocalci...	395	e-109
gi 904210 dbj BAA09453.1  (D50855) Ca-sensing receptor [Homo sa...	395	e-109
gi 1082270 pir  S49341 calcium-sensing receptor - human	395	e-109
gi 7305101 ref NP_038831.1  G protein coupled receptor, family ...	394	e-108
gi 543933 sp P35384 CASR_BOVIN EXTRACELLULAR CALCIUM-SENSING RE...	392	e-108
gi 8393053 ref NP_058692.1  calcium-sensing receptor (hypocalci...	391	e-107
gi 5163340 gb AAD40638.1 AF128842_1 (AF128842) extracellular ca...	391	e-107
gi 1362762 pir  B56715 calcium receptor (clone phPCaR-5.2) - hu...	389	e-107

EST:

	Score	E
gi 3042482 gb AA907022.1 AA907022 oj92a08.s2 Soares_NFL_T_GBC_S...	198	7e-48

**EXPRESSION INFORMATION FOR MODULATORY USE:**

## library source:

Expression information from BLAST EST hit:

gi|3042482|gb|AA907022.1|AA907022 oj92a08.s2 fetal lung NbHL19W, testis NHT, and B-cell (pooled)

Expression information from cDNA library screening:

Human Hela cells  
 Human Bone Marrow

3/10

```

1  MLGPAVLGLS  LWALLHPGTG  APLCLSQQLR  MKGDYVLGGL  FPLGEAEEAG
51  LRSRTRPSSP  VCTRFSSNGL  LWALAMKMAV  EEINNKSDDL  PGLRLGYDLF
101 DTCSEPVVAM  KPILMFILAKA  GSRDIAAYCN  YTQYQPRVLA  VIGPHSSELA
151 MVTGKFFSFF  LMPQVSYGAS  MELLISARETF  PSFFRTVPSD  RVQLTAAAE
201 LQEFGWNNVA  ALGSDDEYGR  QGLSIFSALA  AARGICIAHE  GLVPLPRADD
251 SRLGKVQDVL  HQVNQSSVQV  VLLFASVHAA  HALFNYSISS  RLSPKVWVAS
301 EAWLTSDLVM  GLPGMAQMG  VLGFLOQGAQ  LHEFPQYVKT  HIALATDPAF
351 CSALGEREQG  LEEDVVGQRC  PQDCITLQN  VSAGLNHHQT  FSVYAAVYSV
401 AQALHNTLQC  NASGCPAQDP  VKPWQLLENM  YNLTFHVGG  LPLRFDSSGNV
451 DMEYDLKLWV  WQGSVPRLHD  VGRFNGSLRT  ERLKIRWHTS  DNQKPVSRCS
501 RQCQEQGVRR  VKGFHSCCYD  CVDCEAGSYR  QNPDDIAC  TFCGQDEWSPER
551 STRCFRRRSR  FLAWGEP  AVL  LLLLLLSIAL  GLVLAALGLF  VHHRDSPLVQ
601 ASGGFLACFG  LVCLGLVCLS  VLLFPQGPSP  ARCLAQQPLS  HLP  LPTGCLST
651 LFLQAAEIFV  ESELPLSWAD  RLSGCLRG  FW  AWLVVLLAML  VEVALCTWYL
701 VAFPPPEVTD  WHMLPTEALV  HCRTRSWVSF  GLAHATNATL  AFLCFLGTFL
751 VRSQPGRYNR  ARG  LTFAMLA  YFITWVSEFP  LLANVQVVLR  PAVQMGALLL
801 CVLGILAAFH  LPRCYLLMRQ  PGLNTP  EFFL  GGGPGDAQGQ  NDGNTGNQGK
851 HE

```

(SEQ ID NO:2)

**FEATURES:****Functional domains and key regions:**

[1] PDOC00001 PS00001 ASN\_GLYCOSYLATIONNN-glycosylation site

Number of matches: 9

```

1      85-88  NKSD
2     130-133 NYTQ
3     264-267 NQSS
4     285-288 NYSI
5     380-383 NVSA
6     411-414 NASG
7     432-435 NLTF
8     475-478 NGSL
9     737-740 NATL

```

-----[2]  
 PDOC00004 PS00004 CAMP\_PHOSPHO\_SITEcAMP- and cGMP-dependent protein kinase  
 phosphorylation site  
 556-559 RRRS

-----[3]  
 PDOC00005 PS00005 PKC\_PHOSPHO\_SITEProtein kinase C phosphorylation site  
 Number of matches: 9

```

1     153-155 TGK
2     175-177 SAR
3     189-191 SDR
4     289-291 SSR
5     293-295 SPK
6     477-479 SLR
7     480-482 TER
8     528-530 SYR
9     551-553 STR

```

-----[4]  
 PDOC00006 PS00006 CK2\_PHOSPHO\_SITECasein kinase II phosphorylation site  
 Number of matches: 4

```

1     102-105 TCSE
2     175-178 SARE
3     214-217 SDDE
4     667-670 SWAD

```

-----[5]

4/10

PDOC00008 PS00008 MYRISTYLIN-myristoylation site

Number of matches: 15

1	20-25	GAPLCL
2	69-74	GLLWAL
3	92-97	GLRLGY
4	234-239	GICIAH
5	319-324	GTVLGF
6	476-481	GSLRTE
7	581-586	GLVLAA
8	603-608	GGPLAC
9	646-651	GCLSTL
10	731-736	GLAHAT
11	763-768	GLTFAM
12	804-809	GILAAF
13	831-836	GGGPGD
14	839-844	GQNDGN
15	843-848	GNTGNQ

-----[6]

PDOC00754 PS00980 G\_PROTEIN\_RECEP\_F3\_2G-protein coupled receptors family 3 signature 2  
517-541 CCYDCVDCEAGSYRQNPDDIACATFC

Membrane spanning structure and domains:

Helix	Begin	End	Score	Certainty
1	137	167	1.179	Certain
2	221	241	0.682	Putative
3	256	286	1.149	Certain
4	306	326	1.363	Certain
5	388	408	0.700	Putative
6	567	587	2.340	Certain
7	606	626	2.253	Certain
8	637	657	1.176	Certain
9	680	700	1.695	Certain
10	731	751	1.543	Certain
11	763	783	2.171	Certain
12	792	812	1.955	Certain

5/10

**BLAST Alignment to Top Hit:**

>gi|4337086|gb|AAD18069.1| (AF127389) putative taste receptor TR1  
 [Rattus norvegicus]  
 Length = 840

Score = 460 bits (1170), Expect = e-128

Identities = 276/816 (33%), Positives = 425/816 (51%), Gaps = 39/816 (4%)

```

Query: 31  MKGDYVLGGFLFPL-GEAEEAGLRSTRPSSPVCTR---FSSNGLLWALAMKMAVEEINNKK 86
      + GD++L GLF L G+      L+ R RP      C R      F+ +G      AM+ VEEINN
Sbjct: 33  LPGDFLLAGLFSLHGDC---LQVRHRPLVTSCDRPDSFNHGHYHLFQAMRFTVEEINNS 88

Query: 87  SDLLPGLRLGYDLFDTCSEPVVAMKPSLMFLAKAGSRDIAAYCNYTQYQPRVLAVIGPHS 146
      S LLP + LGY+L+D CSE      + +L LA G R I      + + +V+A IGP +
Sbjct: 89  SALLPNITLGYLELYDVCSESA-NVYATLRVLALQGPRHIEIQKDLRNHSSKVVAFIGPDN 147

Query: 147  SELAMVTGKFFSFFLMPQVSYGASMELLSARETFPSFFRTVPSDRVQLTAAAEELLQEFGW 206
      ++ A+ T      FLMP VSY AS +LSA+ FPSF RTVPSDR Q+      +LLQ FGW
Sbjct: 148  TDHAVTTAALLGPFPLMPLVSYEASSVVLSAKRKFPSFLRTVPSDRHQVEVMVQLLQSFQW 207

Query: 207  NWVAALGSDDEYGRQGLSIFSALAAARGICIAHEGLVPLP-RADDSRLGKVQDVLHQVNQ 265
      W++ +GS +YG+ G+      LA RGIC+A + +VP      R D R+      Q ++ + Q
Sbjct: 208  VWISLIGSYGDYQQLGVQALEELAVPRGICVAFKDIVPFSARVGDPRM---QSMQHLAQ 264

Query: 266  SSVQVLLFASVHAAHALFNYSISSRLSPKVVWVASEAWLTSDLVMGLPGMAQMGTVLGFL 325
      + VV++F++ H A      F      + + L+ KVVWASE W S + + G+ +GTVLG
Sbjct: 265  ARTTVVVVFSNRHLARVFFRSVVLANLTGKVWVASEDWAISTYITSVTGIQIGITVLGVA 324

Query: 326  QRGALHEFPQYVKTHLALAT-----DPAFCSALGEREQGLEEDVVGQRCPCQDCITL 378
      + Q+      ++ ++++ T      + ++CS      Q C +C T
Sbjct: 325  VQQRQVPGLKEFEESYVRAVTAAPSACPEGSWCST-----NQLCRECHTFTT 371

Query: 379  QNVSA--GLNHHQTFSVYAAVYSVAQALHNTLQCNASGCPAQDPVKPWQLLENMYNLTFH 436
      +N+      +      + VY AVY+VA LH L C + C ++ PV PWQLL+ + F
Sbjct: 372  RNMPTLGAFSMSAAYRVYEAVYAVAHGLHQLLGCTSEIC-SRGPVYPWQLLQIQIYKVNFL 430

Query: 437  VGGLPLRFDSNGVNDMEYDLKLWVWQGSVPRRLHDVGRFNGS---LRTERLKIRWHTSDNQ 493
      +      + FD +G+      YD+ W W G      +G + S L      + KI+WH +NQ
Sbjct: 431  LHENTVAFDDNGDTLGYDYDIIAWDWNQPEWTFEIIIGSASLSPVHLDINKTKIQWHGKNNQ 490

Query: 494  KPVSRCRSRQCQEQVRRVKGFHSCCYDCVDCEAGSYRQNPDDIACFTCGQDEWSPERSTR 553
      PVS C+ C G R V G H CC++CV CEAG++      + C CG +EW+P+ ST
Sbjct: 491  VPVSVCTTDCLAGHHRVVVGSHHCCFECVPCCEAGTFLNMSELHICQPCGTEEWAPKESTT 550

Query: 554  CFRRRSRFLAWGEPVILLLLLLLLSLALGLVLAALGLFVHHRDPLVQASGGPLACFGLVC 613
      CF R      FLAW EP L+L+      +L L L++      GLF H +P+V+++GG L L
Sbjct: 551  CFERTVEFLAWHEPISLVLIAANTLLLLLLLVGTAGLFAWHEHTPVVRSAGGRLCFLMLGS 610

Query: 614  LGLVCLSVLLFPQPSPARCLAQQPLSHLPLTGCLSTLFLQAAEIFVESELPLSWADRLS 673
      L      S      F G+P+      CL +QPL L      LS L +++ ++ + +
Sbjct: 611  LVAGSCSFYSFFGEPTVPACLLRQPLFSLGFAIFLSCLTIRSFQLVIIIFKSTKVPTFYR 670

Query: 674  GCLRGPAWLVLVLLAMLVEVALCTWYLVAFPPEVVTDWHMLPTEALVHCRTRSWVSFGLA 733
      + A L V+++ V + +C +LV + P      ++ P      ++ C      + V F LA
Sbjct: 671  TWAQNHGAGLFVIVSSTVHLLICLTWLVMTPTREYQRFPHLVILECTEVNSVGFLLA 730

Query: 734  HATNATLAFLCFLGTFLVRSQPGRYNRARGLTfamLAYFITWVSFVPLLANVQVVLRAV 793
      N L+      F+ ++L + P YN A+ +TF++L F++W++F + + Q PAV
Sbjct: 731  FTHNILLSISTFVCSYLGKELPENYNEAKCVTFSLLLNFVSWIAFFTMAIYQGSYLPVAV 790

```



6/10

Query: 794 QMGALLLCVLGILAAFHLPKCYLLMRQPLNTPEFF 829

+ A L + G + + LP+CY+++ +P LN E F

Sbjct: 791 NVLAGLTTLSGGFSGYFLPKCYVILCRPELNNTTEHF 826

(SEQ ID NO:4)

## Hmmer search results (Pfam):

Model	Description	Score	E-value	N
PF01094	Receptor family ligand binding region	127.0	9.8e-35	2
CE00344	E00344 parathyroid_cell_calcium-sensing_rece	70.5	2.2e-21	4
CE00165	CE00165 Metabotropic glutamate_receptor	58.7	2.1e-18	4
CE00294	E00294 glutamate_receptor_6	26.1	1.4e-07	2
PF00003	7 transmembrane receptor (metabotropic gluta	12.1	0.012	2
PF01796	Domain of unknown function	6.0	2.1	1
PF01160	Vertebrate endogenous opioids neuropeptide	3.6	4.6	1

## Parsed for domains:

Model	Domain	seq-f	seq-t	hmm-f	hmm-t	score	E-value
CE00165	1/4	33	42 ..	1	10 [.	5.8	0.13
CE00344	1/4	32	103 ..	30	102 ..	17.1	4.3e-05
CE00165	2/4	75	104 ..	39	68 ..	9.6	0.008
CE00344	2/4	160	237 ..	161	238 ..	37.4	2.7e-11
CE00294	1/2	75	238 ..	71	264 ..	25.7	1.8e-07
CE00165	3/4	183	239 ..	159	216 ..	29.3	4.5e-09
PF01094	1/2	61	276 ..	1	232 [.	98.3	1.1e-26
PF01796	1/1	352	373 ..	1	23 [.	6.0	2.1
PF01094	2/2	393	456 ..	362	430 ..	28.7	3.3e-07
CE00344	3/4	517	567 ..	562	612 ..	6.2	0.089
CE00294	2/2	396	603 ..	431	649 ..	1.3	3.4
CE00165	4/4	484	603 ..	481	599 ..	13.7	0.00041
PF00003	1/2	566	639 ..	1	72 [.	5.0	1.4
PF01160	1/1	691	709 ..	1	19 [.	3.6	4.6
CE00344	4/4	740	780 ..	784	824 ..	9.8	0.007
PF00003	2/2	740	824 ..	180	270 .]	8.3	0.15

7/10

```

1   GCAAAAGAGG AGAGGCCTGC GTGGGATGTG TGGTGGAGG TGGGGGCCAC
51  TCCAGCCAC  AACAGTGGCC TGGACAGAAA GGGGATGCAA TTCAGCAGAG
101 TCTTCCTGGA TGTACCCCCC ACCTCAGGGT CTGTGGGAGC TGCATATGGT
151 GCCGGCAAAG GCTGCCGACT GCAGTATGGG CCGGGAGAAC TGCCTGGGTC
201 TGGTGGGCC  CCAGGGCAGG GCTCCCTCCG GGTGTTCCTT TCTGTACAAG
251 TGCCATGCTT GTGCCGTTTG CGTGTCCCAA GTGCGAGTGT GCTATTTGCG
301 TGTGCCGCAC GTGTGCCGTT TGCATGTGCT GTTTGCATGT ACCATGTGCA
351 TGTGTGCCAT CTGTGCAATG TGCAGGTGCC AGTTGCATGT GCCATGCGTG
401 TTGGCTGTGA GCGTGTGCTG TTTTCGTGTA TGTGCCATGC ACGTATGTGC
451 TGGTGTGGG  CCGTGCACGT GTGCCACGTG CATGTGTGCC ATTTGTGTAT
501 GCGTGTGTG  CTGTGAGCGT ATGCTGTGCG CATGTGCGTG CCATGCGCTT
551 TGGTGTACC  ATGTGTGTGC TGTTCGATG TGCCATTTGT GTCATGCACG
601 TGCCATTTGC GTGCGATGCG CGTGTGCCAC ACGTCGTTTG CTTGCGTGCC
651 ATGCATGTGT GGCATTTGTG TATGTGCCGT TTCCGTGTGT ATTGTGTGTG
701 CCGTGTGTGT GCCATTTGCA TGTGCCGTTT TGTGTGTGCC ATGCGCGTGT
751 GCCATGCACT TGCCGTGCGT GTGCCATTTG TGTGTACCAT GCGCATGTGC
801 CATTCGTGTG CACCGTACAC GTGTGCCATT TGCATGTATG CTGTGCACGT
851 GCGGCATGCA TGTGTGCCGT TTGCATGCCA TGCATGTGTT CCTTGCGTGT
901 GCGGTGCGTG TCCCATGCAC GTGTGCCGTG CATGTGCCAT TCGCGTGTAC
951 CATGCGCATG TGCCGTTTGT GTGTGTGCGG CATTCTGTG CTCGTGTGCC
1001 AGGTTCGCAT GTGCGCCATA TTCACGTGTG CTCAGCATGT GCCATATGCA
1051 TGTGCGGTGG TAGTGTGTGT CCCTCACAGG TCCTCCTCAC AACACCATGG
1101 GGAAGAAGCA CCAGCCAGGG CACACCTCCT GGTATCTGCT AGGTCTGCCA
1151 GGCCCTAGCT GAAGCTGAGT GCGCCCTCAGT TCCCTGGGA GGGCCTGCGC
1201 CTGGAGTCTG CTGTGTCCCG GAGGGCACCC CCAAAGCAAC ACAGAGGCAG
1251 AGGAGTCCCG GCCCTGCACA CCTGGTGCTG CTCCAGCTGC CGCTCATTTG
1301 CCTGTGGCCC TTCTCCCTT GTTTGCGTGC CCCCCTGGCA AACAACTCT
1351 ACCCCAGCA  GGAGCCACCT GTGTGCCGTC CACGCAGGAG TGGCCAGAC
1401 GGGGGTCAGC AGTGTAGTA  CAGCTGGCCA TGGGTTCTCT ACAGCTTCCA
1451 GGGCTCAGAC TCTGGCAGAA GGGCTGAGAC CCTCAAGGAA CTCTGTCTCC
1501 AAGCAGACTG GGAGGGCAGC ACCACCACCC CAGGGCCCTC CCCAGCTGCA
1551 GGGTGGAGGC CTGGCTGGCC GGCTGCCAC  TGGCCTGACT GGTCTGCAGG
1601 CTTAGGGGGC CCATCCCTGC TGCCCCCGGC TCCGGCCAGC ACAGCCTTGA
1651 GTGGGAGCCA GAAGCTCCCG GGGCTGGGTA GGAGGCATTT CTGTGCTTAT
1701 GAAAAGCCCC AGGGCTGGGT GTCTCTGCAT CCCTCCACG  CAGCTGAGAC
1751 CTCAGAGCCC TGGAGCCCC  TTTGCCCCCT CTCCTCTCCA CAGCCTGCTG
1801 GGCAACTCCA GGAATCGGGG GGTGGCAAGG GGCTCAGCCA CAGGCAGGGA
1851 ACAAGGCCAC GGCCAGCGAC TGAGCAGAGC CTGCCTGCCG GTCAACGCTG
1901 GCCATAGAGC CTGGCAGTGG CCTCAGGCAG AGTCTGACGC GCACAACTT
1951 TCAGGCCCAG GAAGCGAGGA CACCACTGGG GCCCCAGGGT GTGGCAAGTG
2001 AGGATGGCAA GGGTTTGTCT AAACAAATCC TCTGCCCGCT CCCC GCCCG
2051 GGCTCACTCC ATGTGAGGCC CCAGTCGGGG CAGCCACCTG CCGTGCCTGT
2101 TGGAAGTTGC CTCTGCCATG CTGGGCCCTG CTGTCTGGG  CCTCAGCCTC
2151 TGGGCTCTCC TGCACCTGG  GACGGGGGCC CCATTGTGCC TGTACAGCA
2201 ACTTAGATG  AAGGGGACT  ACGTGCTGGG GGGGCTGTTT CCCCTGGGCG
2251 AGGCCAGAGG GGC'TGGCCTC CGCAGCCGGA CACGGCCAG  CAGCCCTGTG
2301 TGCACCAGGT ACAGAGGTGG GACGGCCTGG GTCGGGGTCA GGGTGACCAG
2351 GTCTGGGGTG CTCTGAGCT  GGGGCCGAGG TGGCCATCTG CGGTTCTGTG
2401 TGGCCCCAGG TTCTCCTCAA ACGGCCTGCT CTGGGCACTG GCCATGAAAA
2451 TGGCCGTGGA GGAGATCAAC AACAAAGTCG ATCTGCTGCC CGGGCTGCGC
2501 CTGGGCTACG ACCTCTTTGA TACGTGCTCG GAGCCTGTGG TGGCCATGAA
2551 GCCCAGCCTC ATGTTCTTGG CCAAGGCAGG CAGCCGCGAC ATCGCCGCCT
2601 ACTGCAACTA CACGCAGTAC CAGCCCCGTG TGCTGGCTGT CATCGGGCCC
2651 CACTCGTCAG AGCTCGCCAT GGTCAACGGC AAGTTCTTCA GCTTCTTCCT
2701 CATGCCCCAG GTGGGCGCCC CCCACCATCA CCCACCCCA  CCCAGCCTG
2751 CCCGTGGGAG CCCCTGTGTC AGGAGATGCC TCTTGGCCCT TGCAGGTCAG
2801 CTACGGTGCT AGCATGGAGC TGCTGAGCGC CCGGAGACC  TTCCCTCCT
2851 TCTTCCGCAC CGTGCCAGC  GACCGTGTGC AGCTGACGGC CGCCGCGGAG
2901 CTGCTGCAGG AGTTCGGCTG GAACTGGGTG GCCGCCCTGG GCAGCGACGA
2951 CGAGTACGGC CGGCAGGGCC TGAGCATCTT CTCGGCCCTG GCCGCGCAC
3001 GCGGCATCTG CATCGCGCAC GAGGGCCTGG TGCCGCTGCC CCGTGCCGAT
3051 GACTCGCGGC TGGGGAAGGT GCAGGACGTC CTGCACCAGG TGAACCAGAG
3101 CAGCGTCAG  GTGGTGCTGC TGTTCGCCTC CGTGACGCC  GCCACGCCC
3151 TCTTCAACTA CAGCATCAGC AGCAGGCTCT CGCCCAAGGT GTGGGTGGCC
3201 ACCGAGGCCT GGCTGACCTC TGACCTGTGC ATGGGGCTGC CCGCATGGC
3251 CCAGATGGGC ACGGTGCTTG GCTTCTTCCA GAGGGGTGCC CAGCTGCACG
3301 AGTTCCCCCA GTACGTGAAG ACGCACCTGG CCCTGGCCAC CGACCCGGCC

```

FIGURE 3, page 1 of 4

8/10

```

3351 TTCTGCTCTG CCCTGGGCGA GAGGGAGCAG GGTCTGGAGG AGGACGTGGT
3401 GGGCCAGCGC TGCCCGCAGT GTGACTGCAT CACGCTGCAG AACGTGAGCG
3451 CAGGGCTAAA TCACCACCAG ACGTTCTCTG TCTACGCAGC TGTGTATAGC
3501 GTGGCCCAGG CCCTGCACAA CACTCTTCAG TGCAACGCCT CAGGCTGCCC
3551 CGCGCAGGAC CCCGTGAAGC CCTGGCAGGT GAGCCCGGGA GATGGGGGTG
3601 TGCTGTCCTC TGCAATGTGCC CAGGCCACCA GGCACGGCCA CCACGCCTGA
3651 GCTGGAGGTG GCTGGCGGCT CAGCCCCGTC CCCC GCCCGC AGCTCCTGGA
3701 GAACATGTAC AACCTGACCT TCCACGTGGG CGGGCTGCCG CTGCGGTTCG
3751 ACAGCAGCGG AACCTGGAC ATGGAGTACG ACCTGAAGCT GTGGGTGTGG
3801 CAGGGCTCAG TGCCAGGCT CCACGACGTG GGCAGGTTCA ACGGCAGCCT
3851 CAGGACAGAG CGCCTGAAGA TCCGCTGGCA CACGTCTGAC AACCAGGTGA
3901 GGTGAGGGTG GGTGTGCCAG GCGTGCCCGT GGTAGCCCCC GCGGCAGGGC
3951 GCAGCCTGGG GGTGGGGGCC GTTCCAGTCT CCCGTGGGCA TGCCAGCCG
4001 AGCAGAGCCA GACCCAGGC CTGTGCGCAG AAGCCCCGTG CCCGGTGCTC
4051 GCGGCAGTGC CAGGAGGGCC AGGTGCGCCG GGTCAAGGGG TTCCACTCCT
4101 GCTGCTACGA CTGTGTGGAC TGCAGGCGG GCAGCTACCG GCAAAACCCA
4151 GGTGAGCCGC CTTCGCCGCA GCGGGGGGTG GGAACGCAGC AGGGGAGGGT
4201 CCTGCCAAGT CCTGACTCTG AGACCAGAGC CCACAGGGGA CAAGACGAAC
4251 ACCCAGCGCC CTTCCTCTCT CTCACAGACG ACATCGCCTG CACCTTTTGT
4301 GGCCAGGATG AGTGGTCCCC GGAGCGAAGC ACACGCTGCT TCCGCCGCG
4351 GTCTCGGTTT CTGGCATGGG GCGAGCCGGC TGTGCTGCTG CTGCTCCTGC
4401 TGCTGAGCCT GCGCTGGGC CTGTGTCTGG CTGCTTTGGG GCTGTTCGTT
4451 CACCATCGGG ACAGCCCACT GGTTCAAGGC TCGGGGGGGC CCCTGGCCTG
4501 CTTTGGCCTG GTGTGCCTGG GCCTGGTCTG CCTCAGCGTC CTCCTGTTC
4551 CTGGCCAGCC CAGCCCTGCC CGATGCCTGG CCCAGCAGCC CTTGTCCAC
4601 CTCCCGCTCA CGGGCTGCCT GAGCACACTC TTCTTGCAGG CGGCCGAGAT
4651 CTTCTGTGGG TCAGAACTGC CTCTGAGCTG GGCAGACCGG CTGAGTGGCT
4701 GCCTGCGGGG GCCCTGGGCC TGGCTGGTGG TGCTGCTGGC CATGCTGGTG
4751 GAGGTGCGAC TGTGCACCTG GTACCTGGTG GCCTTCCCGC CGGAGGTGGT
4801 GACGGACTGG CACATGCTGC CCACGGAGGC GCTGGTGAC TGC CGCACAC
4851 GCTCCTGGGT CAGCTTCGGC CTAGCGCACG CCACCAATGC CACGCTGGCC
4901 TTTCTCTGCT TCCTGGGCAC TTTCTTGGTG CGGAGCCAGC CGGGCCGCTA
4951 CAACCGTGCC CGTGGCCTCA CCTTTGCCAT GCTGGCTAC TTCATCACCT
5001 GGGTCTCCTT TGTGCCCCCT CTGGCCAATG TGCAGGTGGT CCTCAGGCCC
5051 GCGGTGCAGA TGGGCGCCCT CTGCTCTGT GTCTTGGGCA TCCTGGCTGC
5101 CTTCCACCTG CCCAGGTGTT ACCTGCTCAT GCGGCAGCCA GGGCTCAACA
5151 CCCCCGAGTT CTTCCTGGGA GGGGGCCCTG GGGATGCCCA AGGCCAGAAT
5201 GACGGGAACA CAGGAAATCA GGGGAACAT GAGTGACCCA ACCCTGTGAT
5251 CTCAGCCCCG GTGAACCCAG ACTTAGCTGC GATCCCCCCC AAGCCAGCAA
5301 TGACCCGTGT CTCGCTACAG AGACCCTCCC GCTCTAGGTT CTGACCCAG
5351 GTTGTCTCCT GACCTGACC CCACAGTAAG CCTAGGCCT GGAGCACGTG
5401 GACACCCCTG TGACCATCTG GGCCCCAGAG CCAAGCTGTG TCCCTGTCCC
5451 TCTGTGCCCA GACCAGGCCT GCCCAGGTAA CCCAGACCCA CTGTTCTGGA
5501 AAGAGGCCCG GAGGGCTCCC AGGGTACCCG CAACCCACAC CGTGAGCTCA
5551 GGAAAAGGAT GCAGGGAGGC CCCGGCCAGA TGGCTGGAAG CCCAAATCAG
5601 GCCCTGCCGA CCTGACCATG TCCCACACAG GCCCCCATCC TGCACCTGC
5651 CAGGCACCAC AGCAGTGGGA GGCCAGGTGG GGGCACACAG GCATATGCCC
5701 AGGGCAGAGC CCGCCGAGGT GGGGGTGGCA CCCAGCTTCC TACTCTGCCC
5751 CTTGCCCAGT GGGTAGACAG CATCATGACT GTCACCAGTA CCAGGGACAG
5801 AGCCGAGTGG GGGTGGGGC GGGGTCCAGC ACCACGGCCA GCACCGACCA
5851 CCAGGACCCC GGAGCCAGCA CCATGGACAG AAAACTGCCC ACCAGGATCT
5901 GACGCCAGCA CGCCGCCAGG CCCACACAGG GTCTCCGGTC AGAGTCCCAG
5951 GGTCACTCC CAGCAGGGCC TAGGGGAGGC TGGACCAGCT CCCTGTGCCT
6001 CATTCCAAGG CAGCCAGCC GGAGAGAAGG GGCACAGGCC ACACATCTGT
6051 CCCATAAAAT TAAACGCTTT TTAGTGTTTA AAATAAGCAG CATTTACACA
6101 GAAGCAGCTC TATGTTAACC ATCTAAACGC TGGGACTTTG ATACAGTATC
6151 TACAGCACAG ACACGTGGGG GCCAGAGAAG CCAGGAAGGC CGCGATGTGT
6201 GCGCGCAGTG TGTGCACTCA CCAAGGACGG GCCACCTGAC TGCCCATCTC
6251 CCCAAGACCT CCCTCCCTGT GGCAGCTGTG CACATCGGGG CCCCTTGACC
6301 CTGCGGGCCA TGGTCCCTCC CTGCCCTGGC TGGGACACGG TGGGCAGGAT
6351 GTCCAGCCCT CTCTCGTCTT CCGGTCCCGT CCTCCACGTA CTTTCAGACT
6401 GTTGCCGGAT GGGAGGAGAG AAGGTGCAGG CTGCTCCAAG GGGCAGCAGC
6451 AGGTGGGACA GATGACAGGG TCGCCTCCTC CCCCAGTCA CCGGCCAGGC
6501 AATAAATAAA TAAATTAGAT CCCTACTCCA GACAGGGGGC CTGTGCACCG
6551 CAGGGGGTTG CCCCCTGGT ACCACCTGG GGGCTGGGC ACAGCTGTGC
6601 AGGAGGTGGG GGTACGCCGA GAGCCCGAGG GGGTCTTCCT CATCCAGGA
6651 GGGATCCCCA CCAGACACAA GGGGTGGGA GGTCCGAGGC TCTCGCTGAG

```

FIGURE 3, page 2 of 4

9/10

```

6701  GGGCAGAGAG  GGAGCGCCCC  CAACACGGCT  GCTCAGACAC  AGGTGCTGTC
6751  AGGAGCTGGA  GCAGCCAGGC  TGCCAGGGCA  GATGGTGGTG  GTCCAGCCTG
6801  CCCCCACCC  TGCTCCCGT  CCTGGCCCCC  ACGAAGGCAA  GCCCACGCGA
6851  GCTCTGCATG  CGGCAGGACC  GCCAGCTCCC  CACCTCAGGC  AGGGCTGGGG
6901  CATGCGCCAC  GAGTCACATG  ATGTCCACGA  AGAACTCGCA  GGGGTTCCCC
6951  ATAGCCTTCT  GGAAGGACTG  GCGGCTGCCT  GTCAATTCCG  GGGGGACGGC
7001  AGCCAGCTCC  CGGACAGGGG  GTCCCCCGGG  TGGCCCCCCC  ACCACTGTAT
7051  AGGCCTTGGT  CGTGGGGTGG  GGCAGGGGGA  GCCCCGGGGC  GGTAGCCGAG
7101  GCCTGACTGC  GTGGGCTGCT  GCCACGGCTG  AGCTGGCCGG  CCGGACGCTC
7151  TCGCCAGCTG  CTCCCCACCC  CACTCGGTGC  CGTGTGATCC  GATTCACTGC
7201  CACTGCCCC  AGTCCCCGCC  GCCCAGCGCT  CCTTCTCACG  GCCCCGGGCC
7251  CGGCGGCTGC  TCCGGGTGGA  CCCACTGCTT  TTGCTCCCTG  GGAGTGAGAA
7301  CAGGATGGGG  AAGGAGCCTG  TCAGCACCAG  GCCCCGGCCAC  GTCCAGAACA
7351  ACCGCCCCCG  CCAGGTGACC  GGGCAGGAAC  GGTGGCCCCG  GCAGGACGGG
7401  TGGGTGGGGC  AGCCATGCAG  GCGCGAGGAC  AGGGCCGGCA  CCCCAGGGT
7451  CGCAGGGGCA  GTCGCGGCAG  GCATGGGCTG  GTGGGGGCAG  TGGGAGGCAG
7501  GCAGGGTCCT  GGGGCAAGCT  GGGCGCCCCC  ACACCTCACC  CCGATGCTTG
7551  AGCTGAGGGC  CTGGTCTTGC  GGCTGAAAGA  CTGAGGTGCC  AGCGAGGGCC
7601  CCTCATGCCC  GGTGCCCCCG  TGGCCATCCT  GGATTCCCCA  CCAAGGCC
7651  CACTGTCCCC  CCGGCCAGG  ACCCTGGCCG  ACGGATGACT  CAGCTCAGCC
7701  CTGTCCTGGG  CTCCAAGAC  GCAGTGGGAG  CTGGAGGGCG  TGGCTGGCTG
7751  GGGACATGCT  GAGGGACCCC  GGGCGGGACC  CTGGCTTACC  GGCCCAAGGT
7801  CCGCTCCGCT  AGTCCTTCAG  TCTAAGGCTT  GTTTAGCACA  AGACAAGGGA
7851  TAGCACGAGT  TACACGCCCC  GCTGCCTGGC  ACCTGCCCCG  CACCCACCCG
7901  CCACCAGTGG  GGA CTGACCG  CGGGCTGGGC  GGGGCTGAAG  TGGGCGCAAG
7951  CGCCGGTCGA  GGGTCAGTAG  ACACCAGCG  TGGCTTCTCT  GCGGTCCAC
8001  A

```

(SEQ ID NO:3)

**FEATURES:**

```

Start:      2118
Exon:       2118-2308
Intron:     2309-2409
Exon:       2410-2710
Intron:     2711-2795
Exon:       2796-3578
Intron:     3579-3692
Exon:       3693-3896
Intron:     3897-4030
Exon:       4031-4151
Intron:     4152-4277
Exon:       4278-5236
Stop:       5234

```

10/10

## MAP POSITION:

#	SHGCNAME	CHROM#	LOD_SCORE	DIST. (cRs)
1	SHGC-57364	1	6.42	32

Bac Accession #: AC026283

## ALLELIC VARIANTS (SNPs):

Position	Major	Minor	Context
406 (SEQ ID NO:5)	t	c	gccatctgtgcaatgtgcaggtgccagttgcatgtgccatgcgtgttggc[t/c] gtgagcgtgtgctgttttcgtgtatgtgccatgcacgtatgtgctgcgtg
852 (SEQ ID NO:6)	t	c	attcgtgtgcaccgtacacgtgtgccatttgcatgtatgctgtgcacgtg[t/c] ggcatgcatgtgtgccgtttgcatgccatgcatgtgttccttgctgtgc
897 (SEQ ID NO:7)	g	a	acgtgcggcatgcatgtgtgccgtttgcatgccatgcatgtgttccttg[c/a] tgtgccgtgcgtgtcccatgcacgtgtgccgtgcatgtgccattcgctg
1,433 (SEQ ID NO:8)	c	t	cgcaggagtggcccagacgggggtcagcagtgtagtacagctggccatg[c/t] ggttcctacagcttcaggcgctcagactctggcagaagggctgagaccct
5,845 (SEQ ID NO:9)	t	c	ggacagagcccaggtgggtgggggcgggtccagcaccacggccagcac[t/c] gaccaccaggaccccgagccagcaccatggacagaaaactgcccaccag
7,028 (SEQ ID NO:10)	g	a	cctgtcaattccgggggacggcagccagctcccgacaggggtcccc[g/a] ggtggccccccaccactgtataggccttggtcgtgggggtggggcggggg

Position	Allele 1	Allele 2	
406	t	c	Intron
852	t	c	Intron
897	g	a	Intron
1,433	c	t	Intron
5,845	t	c	Intron
7,028	g	a	Intron

## SEQUENCE LISTING

<110> Ming-Hui WEI, Wenyan ZHONG, Karen A. KETCHUM, Valentina DIFRANCESCO, Ellen M. BEASLEY

<120> ISOLATED HUMAN G-PROTEIN COUPLED  
RECEPTORS, NUCLEIC ACID MOLECULES ENCODING HUMAN GPCR  
PROTEINS, AND USES THEREOF

<130> CL000869PCT

<160> 10

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 3458

<212> DNA

<213> HUMAN

<400> 1

```

ccccgggtgc aggaattcgg cacgaggcca tgctggggccc tgctgtcctg ggcctcagcc 60
tctgggctct cctgcaccct gggacggggg ccccatgtgt cctgtcacag caacttagga 120
tgaaggggga ctacgtgctg ggggggctgt tccccctggg cgaggccgag gaggctggcc 180
tccgcagccg gacacggccc agcagccctg tgtgcaccag gttctcctca aacggcctgc 240
tctgggcact ggccatgaaa atggccgtgg aggagatcaa caacaagtcg gatctgctgc 300
ccgggctgcg cctgggctac gacctctttg atacgtgctc ggagcctgtg gtggccatga 360
agcccagcct catgttcctg gccaaaggcag gcagcccgca catcgccgcc tactgcaact 420
acacgcagta ccagccccgt gtgttggtgt tcatcgggcc ccactcgtca gagctcgcca 480
tggtcaccgg caagttcttc agcttcttcc tcatgccccca ggtcagctac ggtgctagca 540
tggagctgct gagegcccg gacgaccttc cctccttctt ccgcaccgtg cccagcgacc 600
gtgtgcagct gacggccgcc gggagctgc tgcaggagtt cggctggaac tgggtggccg 660
ccctgggcag cgacgcagag taaggccggc agggcctgag catcttctcg gccctggccg 720
cggcacgcgg catctgcata gcgcacgagg gcctggtgcc gctgccccgt gccgatgact 780
cgcggtggg gaaggtgcag gacgtcctgc accaggtgaa ccagagcagc gtgcaggtgg 840
tgctgctgtt cgcctccgtg cagcccgccc acgccctctt caactacagc atcagcagca 900
ggctctcgcc caaggtgtgg gtggccagcg aggcctggct gacctctgac ctggtcatgg 960
ggctgcccgg catggcccag atggggcacgg tgcttggctt cctccagagg ggtgcccagc 1020
tgcacgagtt cccccagtac gtgaagacgc acctggccct ggccaccgac ccggccttct 1080
gctctgccct gggcgagagg gagcaggggtc tggaggagga cgtggtgggc cagcgctgcc 1140
cgcagtgtag ctgcatacag ctgcagaacg tgagcgagg gctaaatcac caccagagct 1200
tctctgtcta cgcagctgtg tatagcgtgg cccaggccct gcacaacact cttagtgca 1260
acgcctcagg ctgccccgcg caggaccccc tgaagccctg gcagctcctg gagaacatgt 1320
acaacctgac cttccacgtg ggcgggctgc cgctgcggtt cgacagcagc ggaaacgtgg 1380
acatggagta cgacctgaag ctgtgggtgt ggcagggctc agtgcccagg ctccacgacg 1440
tgggcagggt caacggcagc ctacaggacag agcgccgtgaa gatccgctgg cacacgtctg 1500
acaaccagaa gcccggtgtc cgggtgctcg ggcagtgcca ggagggccag gtgcgcgggg 1560
tcaaggggtt ccactcctgc tgctacgact gtgtggactg cgaggcgggc agctaccggc 1620
aaaaccaga cgacatcgcc tgcacctttt gtggccagga tgagtgttcc ccggagcgaa 1680
gcacacgctg cttccgccc aggtctcggt tcttggcatg gggcgagccg gctgtgctgc 1740
tgctgtcctt gctgtgagc ctggcgctgg gccttgtgct ggctgctttg gggctgttgc 1800
ttcaccatcg ggacagccc ctggttcagg cctcgggggg gcccctggcc tgctttggcc 1860
tggtgtgctt gggcctggtc tgccctcagc tcctcctggt ccctggccag cccagccctg 1920
ccgatgcct ggcccagcag cccttgtccc acctcccgt caggggctgc ctgagcacac 1980
tcttctgca ggcggccgag atcttcgtgg agtcagaact gcctctgagc tgggcagacc 2040
ggctgagtggt ctgcctgcgg gggccctggg cctggctggt ggtgctgctg gccatgctgg 2100
tggaggtcgc actgtgcacc tggtaacctg tggccttccc gccggaggtg gtgacggact 2160
ggcacatgct gccacggag gcgctgggtgc actgccgcac acgctcctgg gtcagcttgc 2220
gcctagcgca gccaccaat gccacgctgg ccttctcttg cttcctgggc acttctctgg 2280
tgcgagacca gccggggcgc tacaacctgt cccgtggcct cacctttgcc atgctggcct 2340
acttcatcac ctgggtctcc tttgtgcccc tcctggccaa tgtgcaggtg gtcctcaggc 2400

```

```

ccgccgtgca gatgggcgcc ctccctgctct gtgtccctggg catcctggct gccttccacc 2460
tgcccaggtg ttacctgctc atgcggcagc cagggtctaa caccctcgag ttcttccctgg 2520
gagggggccc tggggatgcc caaggccaga atgacgggaa cacaggaaat caggggaaac 2580
atgagtgacc caaccctgtg atctcagccc cggtgaaccc agacttagct gcgatcccc 2640
ccaagccagc aatgaccctg gtctcgctac agagaccctc ccgctctagg ttctgacccc 2700
aggttgtctc ctgaccctga cccacagtg agccctaggc ctggagcacg tggacacccc 2760
tgtgaccatc tgggccccag agccaagctg tgtccctgtc cctctgtgcc cagaccaggc 2820
ctgcccaggt aaccagacc cactgttctg gaaagaggcc cggagggtc ccagggtacc 2880
cgcaaccac accgtgagct caggaaaagg acgcaggag gccccggcca gatggctgga 2940
agcccaaatc aggcctgcc gacctgacca tgtcccacca gggcccccct cctgcaccct 3000
gccaggcacc acagcagtg gaggcagggt gggggcacac aggcatatgc ccagggcaga 3060
gcccgccgag gtgggggttg caccagctt cctactctgc cccttgcca gtgggtagac 3120
agcatcatga ctgtcaccag taccaggagc agagcccagg tgggggtggg gcggggtcca 3180
gcaccacggc cagcaccgac caccaggacc ccggagccag caccatggac agaaaactgc 3240
ccaccaggat ctgacgccag cagcccgcca ggccacaca gggctctcgg tcagagtccc 3300
agggtcagct cccagcagg cctaggggag gctggaccag ctccctgtgc ctattccaa 3360
ggcagcccag ccggagagaa ggggcacagg ccacacatct gtcccataaa attaaacgct 3420
ttttagtgtt taaaataaaa aaaaaaaaaa aaaaaaaa 3458

```

&lt;210&gt; 2

&lt;211&gt; 852

&lt;212&gt; PRT

&lt;213&gt; HUMAN

&lt;400&gt; 2

```

Met Leu Gly Pro Ala Val Leu Gly Leu Ser Leu Trp Ala Leu Leu His
 1          5          10          15
Pro Gly Thr Gly Ala Pro Leu Cys Leu Ser Gln Gln Leu Arg Met Lys
 20          25          30
Gly Asp Tyr Val Leu Gly Gly Leu Phe Pro Leu Gly Glu Ala Glu Glu
 35          40          45
Ala Gly Leu Arg Ser Arg Thr Arg Pro Ser Ser Pro Val Cys Thr Arg
 50          55          60
Phe Ser Ser Asn Gly Leu Leu Trp Ala Leu Ala Met Lys Met Ala Val
 65          70          75          80
Glu Glu Ile Asn Asn Lys Ser Asp Leu Leu Pro Gly Leu Arg Leu Gly
 85          90          95
Tyr Asp Leu Phe Asp Thr Cys Ser Glu Pro Val Val Ala Met Lys Pro
100          105          110
Ser Leu Met Phe Leu Ala Lys Ala Gly Ser Arg Asp Ile Ala Ala Tyr
115          120          125
Cys Asn Tyr Thr Gln Tyr Gln Pro Arg Val Leu Ala Val Ile Gly Pro
130          135          140
His Ser Ser Glu Leu Ala Met Val Thr Gly Lys Phe Phe Ser Phe Phe
145          150          155          160
Leu Met Pro Gln Val Ser Tyr Gly Ala Ser Met Glu Leu Leu Ser Ala
165          170          175
Arg Glu Thr Phe Pro Ser Phe Phe Arg Thr Val Pro Ser Asp Arg Val
180          185          190
Gln Leu Thr Ala Ala Ala Glu Leu Gln Glu Phe Gly Trp Asn Trp
195          200          205
Val Ala Ala Leu Gly Ser Asp Asp Glu Tyr Gly Arg Gln Gly Leu Ser
210          215          220
Ile Phe Ser Ala Leu Ala Ala Ala Arg Gly Ile Cys Ile Ala His Glu
225          230          235          240
Gly Leu Val Pro Leu Pro Arg Ala Asp Asp Ser Arg Leu Gly Lys Val
245          250          255
Gln Asp Val Leu His Gln Val Asn Gln Ser Ser Val Gln Val Val Leu
260          265          270
Leu Phe Ala Ser Val His Ala Ala His Ala Leu Phe Asn Tyr Ser Ile
275          280          285

```

Ser	Ser	Arg	Leu	Ser	Pro	Lys	Val	Trp	Val	Ala	Ser	Glu	Ala	Trp	Leu
290						295					300				
Thr	Ser	Asp	Leu	Val	Met	Gly	Leu	Pro	Gly	Met	Ala	Gln	Met	Gly	Thr
305					310					315					320
Val	Leu	Gly	Phe	Leu	Gln	Arg	Gly	Ala	Gln	Leu	His	Glu	Phe	Pro	Gln
				325					330					335	
Tyr	Val	Lys	Thr	His	Leu	Ala	Leu	Ala	Thr	Asp	Pro	Ala	Phe	Cys	Ser
			340					345					350		
Ala	Leu	Gly	Glu	Arg	Glu	Gln	Gly	Leu	Glu	Glu	Asp	Val	Val	Gly	Gln
		355					360					365			
Arg	Cys	Pro	Gln	Cys	Asp	Cys	Ile	Thr	Leu	Gln	Asn	Val	Ser	Ala	Gly
	370					375					380				
Leu	Asn	His	His	Gln	Thr	Phe	Ser	Val	Tyr	Ala	Ala	Val	Tyr	Ser	Val
385					390					395					400
Ala	Gln	Ala	Leu	His	Asn	Thr	Leu	Gln	Cys	Asn	Ala	Ser	Gly	Cys	Pro
				405					410					415	
Ala	Gln	Asp	Pro	Val	Lys	Pro	Trp	Gln	Leu	Leu	Glu	Asn	Met	Tyr	Asn
			420					425					430		
Leu	Thr	Phe	His	Val	Gly	Gly	Leu	Pro	Leu	Arg	Phe	Asp	Ser	Ser	Gly
		435					440					445			
Asn	Val	Asp	Met	Glu	Tyr	Asp	Leu	Lys	Leu	Trp	Val	Trp	Gln	Gly	Ser
	450					455					460				
Val	Pro	Arg	Leu	His	Asp	Val	Gly	Arg	Phe	Asn	Gly	Ser	Leu	Arg	Thr
465					470					475					480
Glu	Arg	Leu	Lys	Ile	Arg	Trp	His	Thr	Ser	Asp	Asn	Gln	Lys	Pro	Val
				485					490					495	
Ser	Arg	Cys	Ser	Arg	Gln	Cys	Gln	Glu	Gly	Gln	Val	Arg	Arg	Val	Lys
			500					505					510		
Gly	Phe	His	Ser	Cys	Cys	Tyr	Asp	Cys	Val	Asp	Cys	Glu	Ala	Gly	Ser
		515					520					525			
Tyr	Arg	Gln	Asn	Pro	Asp	Asp	Ile	Ala	Cys	Thr	Phe	Cys	Gly	Gln	Asp
	530					535					540				
Glu	Trp	Ser	Pro	Glu	Arg	Ser	Thr	Arg	Cys	Phe	Arg	Arg	Arg	Ser	Arg
545					550					555					560
Phe	Leu	Ala	Trp	Gly	Glu	Pro	Ala	Val	Leu	Leu	Leu	Leu	Leu	Leu	Leu
				565					570					575	
Ser	Leu	Ala	Leu	Gly	Leu	Val	Leu	Ala	Ala	Leu	Gly	Leu	Phe	Val	His
			580					585					590		
His	Arg	Asp	Ser	Pro	Leu	Val	Gln	Ala	Ser	Gly	Gly	Pro	Leu	Ala	Cys
		595					600					605			
Phe	Gly	Leu	Val	Cys	Leu	Gly	Leu	Val	Cys	Leu	Ser	Val	Leu	Leu	Phe
	610					615					620				
Pro	Gly	Gln	Pro	Ser	Pro	Ala	Arg	Cys	Leu	Ala	Gln	Gln	Pro	Leu	Ser
625					630					635					640
His	Leu	Pro	Leu	Thr	Gly	Cys	Leu	Ser	Thr	Leu	Phe	Leu	Gln	Ala	Ala
				645					650					655	
Glu	Ile	Phe	Val	Glu	Ser	Glu	Leu	Pro	Leu	Ser	Trp	Ala	Asp	Arg	Leu
			660					665					670		
Ser	Gly	Cys	Leu	Arg	Gly	Pro	Trp	Ala	Trp	Leu	Val	Val	Leu	Leu	Ala
		675					680					685			
Met	Leu	Val	Glu	Val	Ala	Leu	Cys	Thr	Trp	Tyr	Leu	Val	Ala	Phe	Pro
	690					695					700				
Pro	Glu	Val	Val	Thr	Asp	Trp	His	Met	Leu	Pro	Thr	Glu	Ala	Leu	Val
705					710					715					720
His	Cys	Arg	Thr	Arg	Ser	Trp	Val	Ser	Phe	Gly	Leu	Ala	His	Ala	Thr
				725					730					735	
Asn	Ala	Thr	Leu	Ala	Phe	Leu	Cys	Phe	Leu	Gly	Thr	Phe	Leu	Val	Arg
			740					745					750		
Ser	Gln	Pro	Gly	Arg	Tyr	Asn	Arg	Ala	Arg	Gly	Leu	Thr	Phe	Ala	Met
		755					760					765			



Leu Ala Tyr Phe Ile Thr Trp Val Ser Phe Val Pro Leu Leu Ala Asn  
 770 775 780  
 Val Gln Val Val Leu Arg Pro Ala Val Gln Met Gly Ala Leu Leu Leu  
 785 790 795 800  
 Cys Val Leu Gly Ile Leu Ala Ala Phe His Leu Pro Arg Cys Tyr Leu  
 805 810 815  
 Leu Met Arg Gln Pro Gly Leu Asn Thr Pro Glu Phe Phe Leu Gly Gly  
 820 825 830  
 Gly Pro Gly Asp Ala Gln Gly Gln Asn Asp Gly Asn Thr Gly Asn Gln  
 835 840 845  
 Gly Lys His Glu  
 850

<210> 3  
 <211> 8001  
 <212> DNA  
 <213> Human

<400> 3  
 gcaaaagagg agaggcctgc gtgggatgtg tgcgtggagg tggggggccac tcccagccac 60  
 aacagtggcc tggacagaaa ggggatgcaa ttcagcagag tcttcctgga tgtcaccccc 120  
 acctcagggt ctgtgggagc tgcataatggt gccggcaaaag gctgccgact gcagtatggg 180  
 ccgggagaaac tgcctgggtc tgcgtgggac ccagggcagg gctccctccg ggtgttgcc 240  
 tctgtacaag tgccatgctt gtgcggtttg cgtgtcccaa gtgcgagtgt gctattttgc 300  
 tgtgccgcac gtgtgccgtt tgcattgtgt gtttgcattg accatgtgca tgtgtgccat 360  
 ctgtgcaatg tgcaggtgcc agttgcatgt gccatgctgt ttggctgtga gcgtgtgctg 420  
 ttttcgtgta tgtgccatgc acgtatgtgc tgcgtgttgg ccgtgcacgt gtgccacgtg 480  
 catgtgtgcc atttgtgtat gccgtgtgtg ctgtgagcgt atgctgtgcg catgtgcgtg 540  
 ccatgcgctt tgcgtgtacc atgtgtgtgc tgtttgcatg tgccatttgt gtcattgcacg 600  
 tgccatttgc gtgcgatgog cgtgtgccac acgtcgtttg cttgcgtgcc atgcatttgt 660  
 ggcattttgt tatgtgccgt ttccgtgtgt attgtgtgtg ccgtgtgtgt gccatttgc 720  
 tgtgccgttt tgtgtgtgcc atgcgcgtgt gccatgcact tgcgtgtcgt gtgccatttg 780  
 tgtgtaccat gcgcattgtgc cattcgtgtg caccgtacac gtgtgccatt tgcattgtatg 840  
 ctgtgcacgt gcggcatgca tgtgtgccgt ttgcatgcca tgcattgtgt ccttgogtgt 900  
 gccgtgcgtg tcccatgcac gtgtgccgtg catgtgccat tcgcgtgtac catgcgcattg 960  
 tgccgtttgt gtgtgtgccc cattcctgtg ctgctgtgcc aggttcgcat gtgcgccata 1020  
 ttcacgtgtg ctcagcatgt gccatattga tgtgcggtgg tagtgtgtgt cctcacagg 1080  
 tcttcctcac aacaccatgg ggaagaagca ccagccaggg cacacctcct ggtatctgct 1140  
 aggtctgcca ggccttagct gaagctgagt gccccccagt tccctggga gggcctgccc 1200  
 ctggagctct ctgtgtcccc gagggcacc ccagagcaac acagaggcag agggatccc 1260  
 gccctgcaca cctgggtgctg ctccagctgc cgtcatttg cctgtggccc ttcctccctt 1320  
 gtttgcgtgc cccctggca aacaaactct acccccagca ggagccacct gtgtgctgc 1380  
 cacgcaggag tggcccagac ggggtcagc agtgtgagta cagctggcca tgcggttcc 1440  
 acagcttcca ggcgtcagac tctggcagaa gggctgagac cctcaaggaa ctctgctccc 1500  
 aagcagactg ggagggcagc accaccaccc caggggccctc cccagctgca ggggtggaggc 1560  
 ctggctggcc ggtgcccac tggcctgact ggtctgcagg cctagggggc ccatccctgc 1620  
 tgccccggc tccggccagc acagccttga gtgggagcca gaagctcccg gggctgggta 1680  
 ggaggcattt ctgtgcttat gaaaagcccc agggctgggt gtctctgcat cctcccacg 1740  
 cagctgagac ctcagagccc tggaggcccc tttgccccct ctcctctcca cagcctgctg 1800  
 ggcaactcca ggaatcgggg ggtggcaagg ggctcagcca caggcaggga acaaggccac 1860  
 ggccagcgac tgagcagagc ctgcctgccg gtcaacgctg gccatagagc ctggcagtgg 1920  
 cctcaggcag agtctgacgc gcacaaactt tcaggcccg gaagcgagga caccactggg 1980  
 gccccagggt gtggcaagt aggatggcaa gggttttgct aaacaaatcc tctgcccgt 2040  
 ccccgccccg ggctcactcc atgtgaggcc ccagtcgggg cagccacctg ccgtgcctgt 2100  
 tggagttgc ctctgccatg ctgggcccct ctgtcctggg cctcagcctc tgggctctcc 2160  
 tgcaccctgg gacggggggc ccatgtgtgc tgtcacagca acttaggatg aagggggact 2220  
 acgtgtctgg ggggtgttcc cccctgggcg aggcgagga ggctggcctc cgcagccgga 2280  
 cacggcccag cagcctgtg tgcaccaggt acagaggtgg gacggcctgg gtcggggta 2340  
 gggtgaccag gtctgggggt ctccctgagct ggggcccagg tggccatctg cggttctgtg 2400  
 tggccccagg ttctcctcaa acggcctgct ctgggcaact gccatgaaaa tggcctgga 2460

ggagatcaac	aacaagtctg	atctgctgcc	cgggctgcgc	ctgggctacg	acctctttga	2520
tacgtgctcg	gagcctgtgg	tggccatgaa	gcccagcctc	atgttccttg	ccaaggcagg	2580
cagccgcgac	atcgccgcct	actgcaacta	cacgcagtac	cagccccgtg	tgctggctgt	2640
catcggggcc	cactcgtcag	agctcgccat	ggtcaccggc	aagttcttca	gcttcttcc	2700
catgccccag	gtgggcgccc	cccaccatca	cccaccccca	cccagccctg	cccgtgggag	2760
ccctgtgtc	aggagatgcc	tcttgccct	tgcaggtcag	ctacggtgct	agcatggagc	2820
tgctgagcgc	ccgggagacc	ttcccctcct	tcttccgcac	cgtgcccagc	gaccgtgtgc	2880
agctgacggc	cgccgcggag	ctgctgcagg	agttcggctg	gaactgggtg	gccgccctgg	2940
gcagcgacga	cgagtacggc	cggcagggcc	tgagcatctt	ctcggccctg	gccgcggcac	3000
gcggcatctg	catcgcgcac	gagggccttg	tgccgctgcc	ccgtgccgat	gactcgcggc	3060
tggggaaggt	gcaggacgtc	ctgcaccagg	tgaaccagag	cagcgtgcag	gtggtgctgc	3120
tgttcgctc	cgtgcacgcc	gcccacgccc	tcttcaacta	cagcatcagc	agcaggctct	3180
cgcccaaggt	gtgggtggcc	agcgaggcct	ggctgacctc	tgacctggtc	atggggctgc	3240
ccggcatggc	ccagatgggc	acgggtgctt	gcttctccca	gaggggtgcc	cagctgcacg	3300
agttccccc	gtacgtgaag	acgcacctgg	ccctggccac	cgaccgggcc	ttctgctctg	3360
ccctgggcga	gagggagcag	ggtctggagg	aggacgtggt	gggccagcgc	tgcccgcagt	3420
gtgactgcat	cacgctgcag	aacgtgagcg	cagggctaaa	tcaccaccag	acgttctctg	3480
tctacgcagc	tgtgtatagc	gtggcccagg	ccctgcacaa	cactcttcag	tgcaacgcct	3540
caggctgccc	cgcgcaggac	cccgtaagc	cctggcaggt	gagcccggga	gatgggggtg	3600
tgctgtcctc	tgcatgtgcc	caggccacca	ggcacggcca	ccacgcctga	gctggagggtg	3660
gctggcggtc	cagccccgtc	ccccgcccgc	agctcctgga	gaacatgtac	aacctgacct	3720
tccacgtggg	cgggctgccc	ctgcggttcg	acagcagcgg	aaacgtggac	atggagtacg	3780
acctgaagct	gtgggtgtgg	cagggctcag	tgcccaggct	ccacgacgtg	ggcaggttca	3840
acggcagcct	caggacagag	cgctgaaga	tccgctggca	cacgtctgac	aaccagggtga	3900
ggtgaggggtg	ggtgtgccag	gcgtgcccg	ggtagccccc	gcggcagggc	gcagcctggg	3960
ggtggggggc	gttccagtct	cccgtgggca	tgcccagccg	agcagagcca	gaccccaggc	4020
ctgtgcgcag	aagcccgtgt	cccgtgctc	gcggcagtcg	caggagggcc	aggtgcgcgc	4080
ggtcaagggg	ttccactcct	gctgctacga	ctgtgtggac	tgcgaggcgg	gcagctaccg	4140
gcaaaaccca	ggtgagccgc	cttcccggca	ggcgggggtg	ggaacgcagc	aggggagggt	4200
cctgccaaagt	cctgactctg	agaccagagc	ccacagggga	caagacgaac	acccagcgcc	4260
cttctcctct	ctcacagacg	acatgcctg	caccttttgt	ggccaggatg	agtggctccc	4320
ggagcgaagc	acacgctgct	tccgcgcag	gtctcggttc	ctggcatggg	gcgagccggc	4380
tgtgctgctg	ctgctcctgc	tgtgagcct	ggcgtggggc	cttgtgctgg	ctgctttggg	4440
gctgttctgt	caccatcggg	acagcccact	ggttcaggcc	tcgggggggc	ccctggcctg	4500
ctttggcctg	gtgtgcctgg	gcctggctct	cctcagcgtc	ctcctgttcc	ctggccagcc	4560
cagccctgcc	cgatgcctgg	cccagcagcc	cttgtcccac	ctcccgtca	cgggctgcct	4620
gagcacactc	ttcctgcagg	cggccgagat	cttcgtggag	tcagaactgc	ctctgagctg	4680
ggcagaccgg	ctgagtggct	gcctgcgggg	gccctggggc	tggctgggtg	tgctgctggc	4740
catgctgggtg	gaggtgcgac	tgtgcacctg	gtacctgggt	gccttcccgc	cggaggtggt	4800
gacggactgg	cacatgcctg	ccacggaggc	gctggtgcac	tgccgcacac	gctcctgggt	4860
cagcttcggc	ctagcgcaag	ccaccaatgc	cacgctggcc	tttctctgct	tcctgggcac	4920
tttctgggtg	cggagccagc	cgggccgcta	caaccgtgcc	cgtggcctca	cttttgccat	4980
gctggccctac	ttcatcacct	gggtctcctt	tgtgcccctc	ctggccaatg	tgaggtggt	5040
cctcaggccc	gccgtgcaga	tgggcgccct	cctgctctgt	gtcctgggca	tcctggctgc	5100
cttccacctg	cccagggtgt	acctgctcat	gcggcagcca	gggctcaaca	cccccgagtt	5160
cttccctggga	gggggcccctg	gggatgccca	aggccagaat	gacgggaaca	caggaaatca	5220
ggggaacat	gagtgaacca	accctgtgat	ctcagccccg	gtgaacccag	acttagctgc	5280
gatccccccc	aagccagcaa	tgaccctgtg	ctcgctacag	agaccctccc	gctctagggt	5340
tgacccccag	gttgtctcct	gaccccgacc	ccacagtaag	ccctaggcct	ggagcacgtg	5400
gacacccctg	tgaccatctg	ggccccagag	ccaagctgtg	tccctgtccc	tctgtgccc	5460
gaccaggcct	gcccaggtaa	cccagaccca	ctgttctgga	aagaggcccg	gagggctccc	5520
agggtagccg	caaccacac	cgtgagctca	ggaaaaggac	gcagggaggc	cccgcccaga	5580
tggctggaag	cccaaatcag	gcctgcccga	cctgaccatg	tcccaccagg	gcccccatcc	5640
tgacccctgc	caggcaccac	agcagtggga	ggccagggtg	gggcacacag	gcatatgcc	5700
agggcagagc	ccgcogaggt	gggggtggca	cccagcttcc	tactctgccc	cttgcccagt	5760
gggtagacag	catcatgact	gtcaccagta	ccagggacag	agcccagggtg	gggtgggggc	5820
ggggctccagc	accacggcca	gcaccgacca	ccaggacccc	ggagccagca	ccatggagac	5880
aaaactgccc	accaggatct	gacgccagca	cgcgcgagg	cccacacagg	gtctccggtc	5940
agagtcccag	ggtcagctcc	cagcagggcc	taggggaggg	tggaccagct	ccctgtgcct	6000
cattccaagg	cagcccagcc	ggagagaagg	ggcacaggcc	acacatctgt	cccataaaat	6060
taaacgcttt	ttagtgttta	aaataagcag	catttacaca	gaagcagctc	tatgttaacc	6120

```

atctaaacgc tgggactttg atacagtatc tacagcacag acacgtgggg gccagagaag 6180
ccaggaaggc cgcgatgtgt gcgcgcagtg tgtgcactca ccaaggacgg gccacctgac 6240
tgcccatctc cccaagacct cctccctgtg ggcagctgtg cacatcgggg ccccttgacc 6300
ctgcgggcca tgggccctcc ctgccctggc tgggacacgg tgggcaggat gtccagccct 6360
ctctcgtctt ccggtcccgt cctccacgta ctttcagact gttgccggat gggaggagag 6420
aaggtgcagg ctgctccaag gggcagcagc aggtgggaca gatgacaggg tcgcctcctc 6480
ccccgagtca ccggccaggc aataaataaa taaattagat ccctactcca gacagggggc 6540
ctgtgcaccg caggggggtt ccccgctggg accaccctgg gggcctgggc acagctgtgc 6600
aggaggtggg ggtcagccga gagcccgagg gggctcttct catcccagga gggatcccca 6660
ccagacacaa ggggttggga ggtccgaggc tctcgtctgag gggcagagag ggagcgcccc 6720
caacacggct gctcagacac aggtgctgtc aggagctgga gcagccaggc tgccagggca 6780
gatggtggtg gtccagcctg cccccaccc tgcctcccgt cctggcccc acgaaggcaa 6840
gccacgcga gctctgcatg cggcaggacc gccagctccc cacctcaggc agggctgggg 6900
catgcgccac gagtcacatg atgtccacga agaactcgca ggggttcccc atagccttct 6960
ggaaggactg gcggctgcct gtcaattccg gggggacggc agccagctcc cggacagggg 7020
gtcccccggt tggccccccc accactgtat aggccttggt cgtgggggtg ggcgggggga 7080
gccccggggc ggtagccgag gcctgactgc gtgggctgct gccacggctg agctggccgg 7140
ccggacgctc tcgccagctg cccccaccc cactcggctg cgtgtgatcc gattcactgc 7200
cactgcccc agctcccgcc gcccgacgct ccttctcacg gcccggggccc cggcggtgc 7260
tccgggtgga cccactgctt ttgctccctg ggagtggaga caggatgggg aaggagcctg 7320
tcagcaccag gcccggccac gtccagaaca accgccccg ccaggtgacc gggcagggaac 7380
ggtggcccggt gcaggacggg tgggtggggc agccatgcag gcgcgaggac agggcgggca 7440
ccccagggt cgcaggggca gctgcggcag gcatgggctg gtgggggagc tgggaggagc 7500
gcagggtcct ggggcaagct gggcgcccc acacctcacc ccgatgcttg agctgagggc 7560
ctggtcttgc ggctgaaaga ctgaggtgcc agcgagggcc cctcatgcc ggtgcccccg 7620
tggccatcct ggattcccca cccaaggccc cactgtcccc ccggcccagg accctggccg 7680
acggatgact cagctcagcc ctgtcctggg ctcccaagac gcagtgggag ctggaggggc 7740
tggctggctg gggacatgct gagggacccc gggcgggacc ctggcttacc ggcccaagg 7800
ccgctccgct agtccttcag tctaaggctt gtttagcaca agacaaggga tagcacgagt 7860
tacagcccc gctgctggc acctgcccgg caccacccc ccaccagtgg ggactgaccg 7920
cgggctgggc ggggctgaag tgggcgcaag cgccggtcga gggtcagtag acaccagcg 7980
tggcttctct gcgtccac a

```

&lt;210&gt; 4

&lt;211&gt; 1138

&lt;212&gt; PRT

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 4

```

Ser Leu Pro Gly Asp Phe Leu Leu Ala Gly Leu Phe Ser Leu His Gly
1          5          10          15
Asp Cys Leu Gln Val Arg His Arg Pro Leu Val Thr Ser Cys Asp Arg
20          25          30
Pro Asp Ser Phe Asn Gly His Gly Tyr His Leu Phe Gln Ala Met Arg
35          40          45
Phe Thr Val Glu Glu Ile Asn Asn Ser Gln Ser Asp Leu Leu Pro Gly
50          55          60
Leu Arg Leu Gly Tyr Asp Leu Phe Asp Thr Cys Ser Glu Pro Val Val
65          70          75          80
Ala Met Lys Pro Ser Leu Met Phe Leu Ala Lys Ala Gly Ser Arg Asp
85          90          95
Ile Ala Ala Tyr Cys Asn Tyr Thr Gln Tyr Gln Pro Arg Val Leu Ala
100         105         110
Val Ile Gly Pro His Ser Ser Leu Leu Pro Leu Gly Tyr Leu Asp Cys
115         120         125
Ser Glu Leu Leu Ala Gly Arg Ile Val Ala Ile Gly Pro Ser Ser Ala
130         135         140
Leu Leu Pro Asn Ile Thr Leu Gly Tyr Glu Leu Tyr Asp Val Cys Ser
145         150         155         160
Glu Ser Ala Asn Val Tyr Ala Thr Leu Arg Val Leu Ala Leu Gln Gly
165         170         175

```

Pro	Arg	His	Ile	Glu	Ile	Gln	Lys	Asp	Leu	Arg	Asn	His	Ser	Ser	Lys
			180					185					190		
Val	Val	Ala	Phe	Ile	Gly	Pro	Asp	Asn	Gln	Ser	Glu	Leu	Ala	Met	Val
		195					200					205			
Thr	Gly	Lys	Phe	Phe	Ser	Phe	Phe	Leu	Met	Pro	Gln	Val	Ser	Tyr	Gly
	210					215					220				
Ala	Ser	Met	Glu	Leu	Leu	Ser	Ala	Arg	Glu	Thr	Phe	Pro	Ser	Phe	Phe
225					230					235					240
Arg	Thr	Val	Pro	Ser	Asp	Arg	Val	Gln	Leu	Thr	Ala	Ala	Ala	Glu	Leu
				245					250					255	
Leu	Gln	Glu	Phe	Gly	Trp	Ala	Thr	Phe	Leu	Met	Pro	Val	Ser	Tyr	Ala
			260					265					270		
Ser	Leu	Ser	Ala	Phe	Pro	Ser	Phe	Arg	Thr	Val	Pro	Ser	Asp	Arg	Gln
		275					280					285			
Leu	Leu	Gln	Phe	Gly	Trp	Ser	Thr	Asp	His	Ala	Val	Thr	Thr	Ala	Ala
	290					295					300				
Leu	Leu	Gly	Pro	Phe	Leu	Met	Pro	Leu	Val	Ser	Tyr	Glu	Ala	Ser	Ser
305					310					315					320
Val	Val	Leu	Ser	Ala	Lys	Arg	Lys	Phe	Pro	Ser	Phe	Leu	Arg	Thr	Val
				325					330					335	
Pro	Ser	Asp	Arg	His	Gln	Val	Glu	Val	Met	Val	Gln	Leu	Leu	Gln	Ser
			340					345					350		
Phe	Gly	Trp	Gln	Asn	Trp	Val	Ala	Ala	Leu	Gly	Ser	Asp	Asp	Glu	Tyr
		355					360					365			
Gly	Arg	Gln	Gly	Leu	Ser	Ile	Phe	Ser	Ala	Leu	Ala	Ala	Ala	Arg	Gly
	370					375					380				
Ile	Cys	Ile	Ala	His	Glu	Gly	Leu	Val	Pro	Leu	Pro	Arg	Ala	Asp	Asp
385					390					395					400
Ser	Arg	Leu	Gly	Lys	Val	Gln	Asp	Val	Leu	His	Gln	Val	Asn	Gln	Trp
				405					410					415	
Gly	Ser	Tyr	Gly	Gly	Leu	Ala	Arg	Gly	Ile	Cys	Ala	Val	Pro	Arg	Asp
			420					425					430		
Arg	Gln	Gln	Ser	Val	Trp	Ile	Ser	Leu	Ile	Gly	Ser	Tyr	Gly	Asp	Tyr
			435				440					445			
Gly	Gln	Leu	Gly	Val	Gln	Ala	Leu	Glu	Glu	Leu	Ala	Val	Pro	Arg	Gly
	450					455					460				
Ile	Cys	Val	Ala	Phe	Lys	Asp	Ile	Val	Pro	Phe	Ser	Ala	Arg	Val	Gly
465					470					475					480
Asp	Pro	Arg	Met	Gln	Ser	Met	Met	Gln	His	Leu	Ala	Gln	Gln	Ser	Ser
				485					490					495	
Val	Gln	Val	Val	Leu	Leu	Phe	Ala	Ser	Val	His	Ala	Ala	His	Ala	Leu
			500					505					510		
Phe	Asn	Tyr	Ser	Ile	Ser	Ser	Arg	Leu	Ser	Pro	Lys	Val	Trp	Val	Ala
		515					520					525			
Ser	Glu	Ala	Trp	Leu	Thr	Ser	Asp	Leu	Val	Met	Gly	Leu	Pro	Gly	Met
	530					535					540				
Ala	Gln	Met	Gly	Thr	Val	Leu	Gly	Phe	Leu	Val	Val	Phe	His	Ala	Phe
545					550					555					560
Leu	Lys	Val	Trp	Val	Ala	Ser	Glu	Trp	Ser	Gly	Gly	Thr	Val	Leu	Gly
				565					570					575	
Ala	Arg	Thr	Thr	Val	Val	Val	Val	Phe	Ser	Asn	Arg	His	Leu	Ala	Arg
			580					585					590		
Val	Phe	Phe	Arg	Ser	Val	Val	Leu	Ala	Asn	Leu	Thr	Gly	Lys	Val	Trp
		595					600					605			
Val	Ala	Ser	Glu	Asp	Trp	Ala	Ile	Ser	Thr	Tyr	Ile	Thr	Ser	Val	Thr
	610					615					620				
Gly	Ile	Gln	Gly	Ile	Gly	Thr	Val	Leu	Gly	Val	Ala	Val	Gln	Gln	Arg
625					630					635					640
Gln	Val	Pro	Gly	Leu	Lys	Glu	Phe	Glu	Glu	Ser	Tyr	Val	Arg	Ala	Val
				645					650					655	

Thr	Ala	Ala	Pro	Ser	Ala	Cys	Pro	Glu	Gly	Ser	Trp	Cys	Ser	Thr	Asn
			660					665					670		
Gln	Leu	Cys	Arg	Glu	Cys	His	Thr	Phe	Thr	Thr	Arg	Asn	Met	Pro	Thr
			675				680					685			
Leu	Gly	Ala	Phe	Ser	Met	Ser	Ala	Ala	Tyr	Arg	Val	Tyr	Glu	Ala	Val
			690			695					700				
Tyr	Ala	Val	Ala	His	Gly	Leu	His	Gln	Leu	Leu	Gly	Cys	Thr	Ser	Glu
705					710					715					720
Ile	Cys	Ser	Arg	Gly	Pro	Val	Tyr	Pro	Trp	Gln	Leu	Leu	Gln	Gln	Ile
				725					730						735
Tyr	Lys	Val	Asn	Phe	Leu	Leu	His	Glu	Asn	Thr	Val	Ala	Phe	Asp	Asp
			740					745					750		
Asn	Gly	Asp	Thr	Leu	Gly	Tyr	Tyr	Asp	Ile	Ile	Ala	Trp	Asp	Trp	Asn
			755				760					765			
Gly	Pro	Glu	Trp	Thr	Phe	Glu	Ile	Ile	Gly	Ser	Ala	Ser	Leu	Ser	Pro
			770			775					780				
Val	His	Leu	Asp	Ile	Asn	Lys	Thr	Lys	Ile	Gln	Trp	His	Gly	Lys	Asn
785					790					795					800
Asn	Gln	Val	Pro	Val	Ser	Val	Cys	Thr	Thr	Asp	Cys	Leu	Ala	Gly	His
				805					810					815	
His	Arg	Val	Val	Val	Gly	Ser	His	His	Cys	Cys	Phe	Glu	Cys	Val	Pro
			820					825					830		
Cys	Glu	Ala	Gly	Thr	Phe	Leu	Asn	Met	Ser	Glu	Leu	His	Ile	Cys	Gln
			835				840					845			
Pro	Cys	Gly	Thr	Glu	Glu	Trp	Ala	Pro	Lys	Glu	Ser	Thr	Thr	Cys	Phe
			850			855					860				
Pro	Arg	Thr	Val	Glu	Phe	Leu	Ala	Trp	His	Glu	Pro	Ile	Ser	Leu	Val
865					870					875					880
Leu	Ile	Ala	Ala	Asn	Thr	Leu	Leu	Leu	Leu	Leu	Leu	Val	Gly	Thr	Ala
				885					890					895	
Gly	Leu	Phe	Ala	Trp	His	Phe	His	Thr	Pro	Val	Val	Arg	Ser	Ala	Gly
			900					905					910		
Gly	Arg	Leu	Cys	Phe	Leu	Met	Leu	Gly	Ser	Leu	Val	Ala	Gly	Ser	Cys
			915				920					925			
Ser	Phe	Tyr	Ser	Phe	Phe	Gly	Glu	Pro	Thr	Val	Pro	Ala	Cys	Leu	Leu
			930			935					940				
Arg	Gln	Pro	Leu	Phe	Ser	Leu	Gly	Phe	Ala	Ile	Phe	Leu	Ser	Cys	Leu
945					950					955					960
Thr	Ile	Arg	Ser	Phe	Gln	Leu	Val	Ile	Ile	Phe	Lys	Phe	Ser	Thr	Lys
				965					970					975	
Val	Pro	Thr	Phe	Tyr	Arg	Thr	Trp	Ala	Gln	Asn	His	Gly	Ala	Gly	Leu
			980					985					990		
Phe	Val	Ile	Val	Ser	Ser	Thr	Val	His	Leu	Leu	Ile	Cys	Leu	Thr	Trp
			995				1000					1005			
Leu	Val	Met	Trp	Thr	Pro	Arg	Pro	Thr	Arg	Glu	Tyr	Gln	Arg	Phe	Pro
			1010			1015						1020			
His	Leu	Val	Ile	Leu	Glu	Cys	Thr	Glu	Val	Asn	Ser	Val	Gly	Phe	Leu
1025					1030					1035					1040
Leu	Ala	Phe	Thr	His	Asn	Ile	Leu	Leu	Ser	Ile	Ser	Thr	Phe	Val	Cys
				1045					1050					1055	
Ser	Tyr	Leu	Gly	Lys	Glu	Leu	Pro	Glu	Asn	Tyr	Asn	Glu	Ala	Lys	Cys
			1060					1065					1070		
Val	Thr	Phe	Ser	Leu	Leu	Leu	Asn	Phe	Val	Ser	Trp	Ile	Ala	Phe	Phe
			1075				1080					1085			
Thr	Met	Ala	Ser	Ile	Tyr	Gln	Gly	Ser	Tyr	Leu	Pro	Ala	Val	Asn	Val
			1090			1095					1100				
Leu	Ala	Gly	Leu	Thr	Thr	Leu	Ser	Gly	Gly	Phe	Ser	Gly	Tyr	Phe	Leu
1105					1110					1115					1120
Pro	Lys	Cys	Tyr	Val	Ile	Leu	Cys	Arg	Pro	Glu	Leu	Asn	Asn	Thr	Glu
				1125					1130						1135

His Phe

<210> 5  
 <211> 101  
 <212> DNA  
 <213> HUMAN

<400> 5  
 gccatctgtg caatgtgcag gtgccagttg catgtgccat gcgtgttggc ygtgagcgtg 60  
 tgctgttttc gtgtatgtgc catgcacgta tgtgctgcgt g 101

<210> 6  
 <211> 101  
 <212> DNA  
 <213> HUMAN

<400> 6  
 attcgtgtgc accgtacacg tgtgccattht gcatgtatgc tgtgcacgtg yggcatgcat 60  
 gtgtgccgtt tgcatgccat gcatgtgttc cttgcgtgtg c 101

<210> 7  
 <211> 101  
 <212> DNA  
 <213> HUMAN

<400> 7  
 acgtgcggca tgcatgtgtg ccgtttgcat gccatgcatg tgttccttgc rtgtgccgtg 60  
 cgtgtcccat gcacgtgtgc cgtgcatgtg ccattcgcgt g 101

<210> 8  
 <211> 101  
 <212> DNA  
 <213> HUMAN

<400> 8  
 cgcaggagtg gccagacgg gggtcagcag tgtgagtaca gctggccatg yggttcctac 60  
 agcttccagg cgtcagactc tggcagaagg gctgagaccc t 101

<210> 9  
 <211> 101  
 <212> DNA  
 <213> HUMAN

<400> 9  
 ggacagagcc caggtggggt gggggcgggg tccagcacca cggccagcac ygaccaccag 60  
 gaccccgag ccagcaccat ggacagaaaa ctgcccacca g 101

<210> 10  
 <211> 101  
 <212> DNA  
 <213> HUMAN

<400> 10  
 cctgtcaatt ccggggggac ggcagccagc tcccggacag ggggtccccc rggtggcccc 60  
 cccaccactg tataggcctt ggtcgtgggg tggggcgggg g 101

## INTERNATIONAL SEARCH REPORT

Int'l Application No

PCT/US 01/07832

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/705 C07K16/28 C12N15/12 A01K67/02 C12P21/00  
C12Q1/68 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, SEQUENCE SEARCH

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 01 64882 A (MILLENNIUM PHARMACEUTICALS) 7 September 2001 (2001-09-07) the whole document, especially figure 15 (seq.id. no. 13 and 15)	1-16, 19-23
X	<p>-----</p> <p>DATABASE EM_HTG 'Online! embl; 13 February 2000 (2000-02-13) LEONGAMONLERT, D.: "Human DNA from clone RP5-89003" retrieved from EBI, accession no. AL139287 Database accession no. AL139287 XP002182829 99,9% identity in 1814 nt overlap (1625-3438:69303-71116) abstract</p> <p>-----</p> <p style="text-align: center;">-/--</p>	4-6,8,9



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* & \* document member of the same patent family

Date of the actual completion of the international search

15 November 2001

Date of mailing of the international search report

28/11/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

De Kok, A

## INTERNATIONAL SEARCH REPORT

Int onal Application No

PCT/US 01/07832

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EM_EST 'Online! EMBL; 12 July 1999 (1999-07-12) STRAUSBERG, R.: "Human cDNA clone IMAGE:2419799 3' end" retrieved from EBI, accession no. AI818485 Database accession no. AI818485 XP002182830 94% identity in 411 nt overlap (3458-3048:15-425) abstract ----	4-6,8,9
A	WO 00 06592 A (UNIVERSITY OF CALIFORNIA) 10 February 2000 (2000-02-10) the whole document, especially pages 79-80 and claim 5 ----	1-16, 19-23
A	US 5 763 569 A (BROWN EDWARD M ET AL) 9 June 1998 (1998-06-09) the whole document, especially figure 47 ----	1-16, 19-23
A	HOON M A ET AL: "PUTATIVE MAMMALIAN TASTE RECEPTORS: A CLASS OF TASTE-SPECIFIC GPCRS WITH DISTINCT TOPOGRAPHIC SELECTIVITY" CELL, MIT PRESS, CAMBRIDGE, MA,, US, vol. 96, 19 February 1999 (1999-02-19), pages 541-551, XP000922524 ISSN: 0092-8674 the whole document -----	1-16, 19-23



# INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/US 01/07832

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0164882	A	07-09-2001	WO 0164882 A2	07-09-2001
WO 0006592	A	10-02-2000	AU 5238199 A	21-02-2000
			BR 9912545 A	30-10-2001
			CN 1317010 T	10-10-2001
			EP 1100810 A1	23-05-2001
			NO 20010363 A	26-03-2001
			WO 0006592 A1	10-02-2000
US 5763569	A	09-06-1998	US 6011068 A	04-01-2000
			US 5962314 A	05-10-1999
			AU 709303 B2	26-08-1999
			AU 4195796 A	15-05-1996
			BR 9509411 A	30-12-1997
			CA 2202879 A1	02-05-1996
			CN 1220658 A	23-06-1999
			CZ 9701182 A3	17-09-1997
			EP 0787122 A2	06-08-1997
			HU 77980 A2	01-02-1999
			JP 11130737 A	18-05-1999
			JP 2882882 B2	12-04-1999
			JP 10513436 T	22-12-1998
			NZ 297157 A	30-08-1999
			PL 319812 A1	01-09-1997
			WO 9612697 A2	02-05-1996
			US 6001884 A	14-12-1999
			US 5858684 A	12-01-1999
			US 6031003 A	29-02-2000
			US 5688938 A	18-11-1997
			US 6211244 B1	03-04-2001
			AU 3122699 A	22-07-1999
			AU 702629 B2	25-02-1999
			AU 8087294 A	08-05-1995
			CA 2173747 A1	27-04-1995
			CN 1139917 A	08-01-1997
			EP 0724561 A1	07-08-1996
			JP 9504032 T	22-04-1997
			WO 9511221 A1	27-04-1995
			SG 52796 A1	28-09-1998
			WO 9304373 A1	04-03-1993
			AU 673500 B2	14-11-1996
			AU 711247 B2	07-10-1999
			AU 7197796 A	20-02-1997
			CA 2115828 A1	04-03-1993
			CN 1071333 A ,B	28-04-1993
			EP 0657029 A1	14-06-1995
			IL 102917 A	06-12-2000
			JP 2860285 B2	24-02-1999
			JP 9328420 A	22-12-1997
			JP 2887201 B2	26-04-1999
			JP 9281109 A	31-10-1997
			JP 11221095 A	17-08-1999
			JP 2728564 B2	18-03-1998
			JP 6510531 T	24-11-1994
			JP 2001220356 A	14-08-2001
			KR 251995 B1	01-06-2000
			MX 9204881 A1	01-04-1993
			NO 940581 A	25-04-1994

# INTERNATIONAL SEARCH REPORT

### Information on patent family members

Internal Application No

PC1/US 01/07832

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5763569	A	RU 2147574 C1	20-04-2000